
RESPONSE LETTER

-- Ref1.1 – Significance about the results of MET --

Reviewer Comment	The authors have focused on MET and produced some data that did not provide further advances to what we have known so far on the role of MET in type I pRCC.
Author Response	<p>We thank the reviewer for expressing concerns about our results on MET. Indeed, MET has been known to be the central driver in type I pRCC for decades. However, most of the analyses focus on coding region only. The majority of type I pRCC patients in the TCGA study do not carry any missense mutation in MET. ~20% patients show significantly higher MET expression yet are completely silent in MET, without evidence for missense mutations, alternative splicing and copy number amplification. Using a more integrated approach, particularly focusing on non-coding, we are able to provide hints for alternative mechanisms to MET dysfunction in type I pRCC. <u>In particular:</u></p> <ol style="list-style-type: none">1. Our study is the first one that looks into the non-coding regions of pRCC. It is an open question in the field of cancer genomics, whether whole genome sequencing adds additional value over exome sequencing. Recent studies in whole genome sequencing suggest active roles of non-coding mutations in cancer. Well-known examples include TERT promoter mutations in urothelial carcinoma and enhancer hijacking in CNS tumors (REF). However, the debate of WGS versus WES remains unsettled and significances of many non-coding alterations stay unknown. In this study, we investigate the functional roles of non-coding alterations in pRCC. We find excessive non-coding mutations at the promoter and regulatory regions of MET. Given the critical role MET plays in pRCC and some MET-driven samples are completely silent in terms of alterations of MET, we believe this mutation hotspot is possibly linked with pRCC molecular etiologies. Accordingly, we have revised the manuscript to better explain the significance of our findings.2. During our revision, we find the activation of a cryptic promoter in the second intron of MET causes the alternative mRNA isoform described in the original TCGA study. This event has been observed in several other cancers included CML and some GI (gastrointestinal tract) cancers. We provide an explanation for the alternative MET transcription isoform in pRCC. Further more, we

	<p>linked the usage of this cryptic promoter with the methylation change that is often seen in pRCC. We added this new analysis in the revised manuscript.</p> <p>3. We find more somatic mutations in an extended WXS set, further completing the MET mutation spectrum of the TCGA study.</p>
Excerpt From Revised Manuscript	<p>The TCGA study identified a MET alternative translation isoform as a driver event (3). However, the etiology of this new isoform was unknown. We found this isoform results from the usage of a cryptic promoter from an L1 element, likely due to local loss of methylation (REF). This event was reported in several other cancer types (REF). To test its relationship with methylation, we found the closest probe (cg06985664, ~3kb downstream) on the Methylation array show marginally statistically significant ($p=0.055$, one-side rank-sum test). Additionally, as expected, this event is associated with methylation group 1 (odds ratio (OR)= 4.54, 95%CI: 1.07-19.34, $p<0.041$), indicating genome-wide methylation dysfunction. This association is stronger in type 2 pRCC and it shows a significant association with the C2b cluster (OR= 17.5, 95%CI: 1.72-32.6, $p<0.007$).</p>

-- Ref1.2 – Non-coding analysis power--

Reviewer Comment	<p>The non-coding analysis did not show significant findings, likely due to the small cohort size and the heterogeneous nature (cohort (n=32) included 19 type I pRCC, 6 type II pRCC, and 7 unclassified).</p>
Author Response	<p>We agree with the referee that our statistical test power is affected. However, in terms of the size of data, our 35 WGS samples (with three newly added samples, see below) have more reads than >1,000 WES samples. Unlike the traditional statistics test by looking at a single (or a few) feature in the cohorts, we have the ability to obtain a large number of features (in forms of genomic regions) about each sample. This forms a high-dimensional scenario ($p \gg n$, "short, fat data") commonly seen in big data analysis nowadays. As the referee points out, low cohort number limits our statistical power. But we instead conduct a comprehensive and unbiased examination of the entire genome for each sample. The great amount of data we acquire from every single sample greatly boosts our analyses. The impactful non-coding alterations we identify are in fact filtered out from thousands of changes in the entire genome and we have high confidence that they have truly high impacts.</p> <p>In our study, we are able to show a significant amount of samples carry impactful mutations in noncoding regions and conduct some coarse recurrence tests. Our analysis is the first exploration of pRCC non-coding regions and provides meaningful insights of pRCC. This hopefully will spark some research ideas and interests in noncoding regions of pRCC.</p> <ol style="list-style-type: none"> 1. The non-coding mutation hot spots indeed carry excessive and impactful mutations. We segment the genome based on functional annotation (FunSeq). Then we try to find

	<p>highly recurrent mutations in annotated regions. These three mutation hotspots have extremely high mutation rate in our cohort. The hotspots span from 7 to 50kb, each with 6-to-7 mutations observed in 35 samples (~150,000 non-coding mutations in total). Therefore, the local mutation rate is roughly 5-to-20 times higher than average. We explain our approaches better in the revised manuscript.</p> <ol style="list-style-type: none"> 2. We leverage the existing knowledge in coding regions and complete the picture of cancer genomes with our non-coding analyses. All three hotspots are tightly linked with coding genes that are biologically associated with pRCC. Mutations in these regions could have high impacts. Unfortunately, non-coding regions are largely overlooked in the previous studies of pRCC. Our study is the first one that looks into these regions that make up to 98% of the genome. Although we were not able to perform fine-scale tests for these mutation hotspots due to sample size, we hope our analyses will spark interests and encourage researcher to further explore the possible biological impacts of these events. 3. In our revision process, we reviewed the WGS samples and added three more WGS samples into our cohort, reaching a final size of 35. We also want to point out that, because WGS covers more than 50 times more regions than WES, additional three samples add more reads than 100 WES samples. Thus this added data greatly boosted our data-driven analyses, for example, signature and mutation landscape study.
Excerpt From Revised Manuscript	

Shantao 2/12/2017 6:36 AM
Deleted: ~

-- Ref1.3 – Implications of NEAT1 mutations--

Reviewer Comment	This reviewer was very intrigued by the NEAT1 finding, which deserves more work to elucidate its importance and could be the highlight of this paper. Can we use NETA1 promoter mutation to classify pRCC and what are the associated transcriptomic signature?
Author Response	<p>Recurrent mutations in NEAT1 are indeed intriguing. NEAT1 is a non-coding RNA thus will be missed by whole exome sequence. It was overlooked in previous studies of pRCC. Our study is the first one on NEAT1 in pRCC. We show a mutation hotspot in NEAT1 and mutations are linked with higher expression of NEAT1, presumably due to the dysfunction of gene regulation region, and worse survival of patients. As the referee suggested, we did additional work on NEAT1 in the revised manuscript.</p> <p>Although lacking WGS data to find genomic alteration, we found</p>

	<p>NEAT1 is overexpressed in 5% of the TCGA ccRCC cohort. NEAT1 higher expression is significantly associated with shorter overall survival time (median OS: 36 months versus 77 months). NEAT1 is tightly co-expressed with MALAT1 in both pRCC and ccRCC, which is another noticeable lncRNA in cancer.</p> <p>The referee raised an interesting point about expression signature. NEAT1 mutations seem to be associated with RNAseq cluster 3 but do not reach statistical significance (p>0.05), likely due to small sample size. NEAT1 expression pattern is...</p> <p>We expect with a larger cohort, the statistical significance we get will be strengthened. As an active participant of the currently ongoing PCAWG (PanCancer Analysis of Whole Genomes), we quickly look into NEAT1 mutations in the high quality PCAWG RCC dataset. 21/144(14.58%) of the samples carry mutations in NEAT1, a frequency agrees with the one from our cohort.</p> <p>WE add a new section and a supplemental figure to reflect the discussions above and our new analysis results of NEAT1. In particular, we emphasize that, with ...</p>
Excerpt From Revised Manuscript	<p>However, without mutation status, NEAT1 expression level is not significantly linked with pRCC survival. Nonetheless, NEAT1 is overexpressed in 5% ccRCC samples from the TCGA cohort. NEAT1 overexpression is significantly associated with shorter overall survival (Fig SXX). MALAT1, another noticeable lncRNA in cancer, is tightly co-expressed with NEAT1 in both pRCC and ccRCC. Overexpression of MALAT1 is reported to be associated with cancer progression (REF).</p>

Shantao 2/12/2017 6:39 AM
Deleted: about 6

Shantao 2/12/2017 6:38 AM
Deleted: study

Shantao 2/12/2017 6:40 AM
Deleted: are allured

Shantao 2/12/2017 6:40 AM
Deleted: to

Shantao 2/12/2017 6:38 AM
Deleted: Unfortunately, we are not able to publish results based on PCAWG data at this moment.

-- Ref1.4 –Significance of mutation spectra & landscape--

Reviewer Comment	<p>The findings on mutation spectra and defects in chromatin remodeling affecting mutation landscape are of moderate interest.</p>
Author Response	<p>We appreciate the referee for raising concerns about the significance of the mutation spectra and landscape analysis.</p> <p>Several recent landmark pan-cancer studies lead to the wide recognition of significance and great research interests in cancer mutational processes (REF). DNA mutation is one of the important driving forces of cancer development. Understanding the underlying processes and affecting factors that generate the mutations is vital in cancer studies.</p> <p>As the referee points out earlier, pRCC is very heterogeneous, especially the type II. TCGA study shows several subgroups of pRCC while we still observe great variation in subgroups. A key aim of our study is to better understand this heterogeneity. Studying the heterogeneity in the fundamental processes that generate the mutations brings about great research excitement.</p>

	<p>Mutation spectra elucidate diversified mutation processes in pRCC. In our study, we identify several factors (methylation, APOBEC, chromatin remodeling defects etc.) <u>that</u> play important roles in tumorigenesis. This helps better characterize and understand pRCC in terms of variations in mutagenesis, tumor evolution, and molecular etiologies. It also has potential clinical implications. For instance, mutation burden has important predictive value on immune therapy response. In the era of great advancing of immune therapy, we feel research on mutation landscape in pRCC has the potential to facilitate clinical decisions.</p> <p><u>Moreover, WGS provides unique advantages to mutation spectra and landscape analyses. First, WGS produces more than an order more SNVs in the sample, which greatly powers the analyses. Second, most of the noncoding regions are not subject to strong selection and thus minimizes the selection confounding effects. Last, WGS does not need the exon capture preprocess step before sequencing. Therefore, WGS avoids related bias and technical artifacts.</u></p> <p>In the revised manuscript we add discussions to better explain the significance of this part of the study.</p>
Excerpt From Revised Manuscript	

-- Ref1.5 – Individual evolution trees --

Reviewer Comment	The WGS analysis is somewhat descriptive. With the wealth of this dataset, the author shall attempt to generate individual pRCC evolution trees of these 32 cases.
Author Response	<p>We thank the referee for the suggestion. <u>WGS indeed provides a large amount of SNVs (> 16,000 in total, which equals more than 100 WES samples), making our dataset suitable for evolution analysis.</u> In the revision, we build <u>individual evolution trees for 35 cases.</u></p> <p><u>We identify four major types of trees, based on topology and the number of subclones identified: linear (no branch) with fewer subclones, linear with more subclones, shortly branched and branched with distal subclones. We find some interesting correlations of tree topology and cancer subtypes. Trees with short branches are enriched in type 1 RCC (p<0.002) while trees with long branches or linear tree with more subclones occurs</u></p>

	<p>more often in type 2. The evolution tree structure might reflect some intriguing cancer biological processes in RCC. Type I RCCs are enriched in more homogenous tree structures. Subclones exist but they are not divergent. Whereas type II RCCs have more subclones and show greater degree of divergence. This is in line with the knowledge that type II RCCs are more heterogeneous.</p> <p>We add a new figure panel in figure 4 and corresponding results/discussion sections for the new evolution tree analysis. Also, we show 35 individual trees, one for every single sample, in the supplement figures.</p>
Excerpt From Revised Manuscript	

-- Ref1.5 – Minor --

Reviewer Comment	<p>a) line 173, please add reference b) line 258, based on available clinical trials, there is almost certain that c-MET inhibitor has no role in type II pRCC, which needs to be rephrased. c) line 278-283, will expand pending further analysis</p>
Author Response	<p>We thank the referee for pointing out these issues. In the revision, we</p> <p>a) added reference to support higher mutation rate of C-to-T in methylated CpGs. (T.R. Waters, P.F. Swann Thymine-DNA glycosylase and G to A transition mutations at CpG sites Mutat. Res., 462 (2000), pp. 137–147) b) “Potentially, patients with rs11762213 might also benefit from MET inhibitors Should we fight back on this? Stating “Type II patients carrying rs11762213 only constitute a small subset of the patients. Thus clinical trials were not able to rule out MET inhibitor might be effective in this subset. One following-up study could be stratifying the cohort based on rs11762213 genotype and reanalyze the data”</p> <p>I think Brian has some ongoing MET inhibitor trials to support this. Or we just turn down the language? c) We expanded the section of NEAT1. See REF1.3.</p>
Excerpt From Revised Manuscript	b)



-- Ref2.1 – Molecular mechanisms of rs11762213 --

Reviewer Comment	For the germline SNP rs11762213, it does not change protein sequence. If it really plays some role in cancer, it probably has regulatory function(s). However, the authors didn't observe changes in expression or protein abundance of MET. I am wondering what about the expression and protein abundance of MET in ccRCC where this SNP also is associated with prognosis. And what about genes that are next to MET in both pRCC and ccRCC if MET is unchanged?
Author Response	<p>The referee raised an excellent question. The mechanism of rs11762213, a synonymous exonic SNP, remains still unsettled. A recent publication about rs11762213 by AA Hakimi et al. studies this in great details in ccRCC. They did not find any statistically significant change in MET expression patterns associated with this SNP. Also this SNP is not in strong linkage disequilibrium with other SNPs of interest in RCCs.</p> <p>Following the suggestion of referee, we explored the genes within 50kb away from MET ...</p> <p>Since this is a germline SNP, it may affect the tumor development, even at the very early stage. Such effects might be complicated and become cryptic during the tumor development and thus fail to be detected. Also this SNP might have affect the MET expression in nearby tissues and stimulate the tumor growth. AA Hakimi et al., were not able to get statistical significance on higher MET expression in normal tissue associated with rs11762213. However, this could be due to low statistical power.</p> <p>In the revised manuscript, we better elaborate the current research status of rs11762213 and incorporate the discussions above.</p>
Excerpt From Revised Manuscript	

-- Ref2.2 – DHS validity --

Reviewer Comment	The authors shall use caution when counting mutations in DHS sites when there is mutation in chromatin remodelers. The authors claimed mutations in chromatin remodelers can change the chromatin environment. If so, comparing number of mutations in DHS sites predicted from one cell line will particularly be problematic in patients with mutations in remodelers.
Author Response	The referee made an excellent observation. We certainly agree that, DHS regions called from a normal kidney cell line represent the open chromatin regions under normal, physiological condition. With chromatin remodeling dysfunction, the DHS regions are likely to shift in pRCC tumors. In fact we believe this

<p style="color: red; font-size: 2em; transform: rotate(-45deg);">FURTHER DIFF</p> <p style="color: red; font-size: 1.5em; transform: rotate(-45deg);">DOES IT MAKE A DIFF</p>	<p>is a very plausible explanation for mutation landscape changes since chromatin environment greatly affects DNA repair and replication. We admit the language we use in the manuscript causes confusion. In the revised manuscript, we use “open chromatin regions in normal kidney cells” to accurately describe the nature of these DHS sites.</p> <p style="background-color: yellow;">In the revision, we substitute the cell line with eleven fetal kidney cortex tissues. Using kidney tissues better reflects the physiological, normal kidney tissue. Also gathering data from eleven tissues allows us minimizing experimental bias and variance. Last, since the eleven tissues come from different fetus with different genetic background, we are able to reduce individual variance and call conservative/stable DHS sites by taking the intersection of the DHS regions.</p> <p>Last, DHS regions are enriched with functional regions of genome, for example, essential genes. Therefore, a higher mutation burden in DHS regions might be deleterious for tumor. Nonsynonymous mutations in protein coding regions may also be antigenic. Recent studies have shown patients with higher and impactful mutation burden response better to immunotherapy. Thus this shift of mutation landscape may have clinical implications.</p>
Excerpt From Revised Manuscript	

Shantao 2/12/2017 7:37 AM
Formatted: Highlight

-- Ref2.3 – Figure 2A --

Reviewer Comment	Figure 2A is confusing. There are 3 proposed promoters and 4 SNVs in promoter, inconsistent with text. It's better to put this panel into Figure 1 rather than in Figure 2.
Author Response	We thank the reviewer for pointing the flaws in our figure preparation. We have fixed the promoter regions and put it into Figure 1.
Excerpt From Revised Manuscript	

-- Ref2.4 – Color key in Figure 4 --

Reviewer Comment	Color key should be added in Figure 4
Author Response	We thank the reviewer for pointing the flaws in our figure preparation. We have added color key in Figure 4
Excerpt From Revised Manuscript	

-- Ref3.1 – The significance of rs11762213 in pRCC--

Reviewer Comment	They looked at an exonic SNP in the MET gene among pure papillary RCC (rather than mixed RCC histologies done previously) and found marginally worse prognosis in type 2 pap RCC with the SNP. They argue that this may have clinical implications and that patients with the SNP may benefit from MET inhibitors. However, the association is not strong enough for it to matter clinically. A cost benefit analysis would be needed as well as an explanation of how it would impact management. The claim that it would select patients for MET inhibition is unsubstantiated. The authors link this SNP to a racial predisposition to developing papillary RCC ... but this is mostly speculation.
Author Response	<p>We totally agree with the reviewer that there is a long path to translate scientific discoveries in the lab into clinical care. In this scientific research article, we have no intention to offer any suggestion for clinical practice changes. Cost-benefit analysis and many more studies are certainly needed before any change in patient management. We are afraid that they are beyond the scope of the article and <i>Plos Genetics</i>.</p> <ol style="list-style-type: none"> 1. The two previous studies about rs11762213 were done on a mixed RCC cohort and a cohort entirely made up of ccRCC respectively. The mixed cohort was mostly ccRCC (78% in discovery cohort and 75% in validation cohort) due to the disease nature. The pRCC subset is apparently too small to run any subgroup analysis. Both of the studies were not able to prove rs11762213 predict prognosis in pRCC. In this manuscript, for the first time, we find that rs11762213 has predictive value in type 2 pRCC outcome. 2. p-value indicates the chances that the null hypothesis is true. It is certainly impacted by the magnificence of the effects of the SNP. But, many other factors also greatly affect the p-value, for example, statistical power/sensitivity. In our case, the p-value is largely bounded by the small sample size. A “marginal” p-value does not necessarily mean the effect of the SNP on prognosis is small. In the revised manuscript, we calculated the odds ratio to better reflect the effect of rs11762213. 3. We were forming hypotheses and speculating about the etiologies and implications of rs11762213 in the discussion section. <p>We agree with the reviewer that we should rewrite this part to better explain the implications of our study. Thus we revised the SNPs discussion in the manuscript.</p>
Excerpt From Revised Manuscript	

-- Ref3.2 –Statistical significance--

Reviewer Comment	Their analysis of non-coding mutation hotspots was largely negative or statistically underpowered. They found mutations in the promoter region of NEAT1, a non-coding RNA, which were marginally associated with worse outcome. This is interesting but of minor significance.
Author Response	We understand the concern of the reviewer. However, we feel the recurrent mutations in NEAT1 are actually of great interest. First, NEAT1 is a non-coding RNA thus will be missed by whole exome sequence. It was overlooked in previous studies of pRCC. We conducted the first study of NEAT1 in pRCC. ...same as REF1.3
Excerpt From Revised Manuscript	"However, without mutation status, NEAT1 expression level is not significantly linked with pRCC survival. Nonetheless, NEAT1 is overexpressed in about 6% ccRCC samples from the TCGA cohort. NEAT1 overexpression is significantly associated with shorter overall survival (Fig SXX). MALAT1, another noticeable lncRNA in cancer, is tightly co-expressed with NEAT1 in both pRCC and ccRCC. Overexpression of MALAT1 is reported to be associated with cancer progression (REF)."

-- Ref3.3 – Interpretation of APOBEC--

Reviewer Comment	They found an APOBEC mutation signature in only 1 out of 155 cases. Given that APOBEC signatures are described in urothelial carcinoma, the authors then theorized that papillary RCC may be genomically similar to urothelial carcinoma ... and may potentially be managed similarly with chemotherapy and radiation therapy. This is a great leap of faith and logic (or illogic). Again, attesting to the paucity of actual positive findings.
Author Response	We thank the reviewer for expressing the concerns about our interpretation of APOBEC and the language we use here. pRCC is very heterogeneous, especially the type II. TCGA study shows several subgroups of pRCC and still we see large variation within subgroups. A key aim of our study is to better understand this heterogeneity. APOBEC mutagenesis shows both location (prefer single-strand DNA, for example around double strand break sites) and context (unique trinucleotide signature) preference. Therefore, in APOBEC active samples, it is a major player in shaping the cancer genome. In previous clinical studies, ~15% of pRCC patients response to cytotoxic chemo (REF) but we do not know who they are. Our APOBEC study and comparison to urothelial cancer are making efforts to better understand the heterogeneity of the cancer nature. We want to emphasize that we are now doing explorations and forming hypotheses, trying to raise further research interests.

	<p>We were forming scientific hypotheses here in the discussion section in hope to encourage further research ideas and interests. We completely understand the concern from the reviewer about the language and interpretation of the results. Therefore, in the revised manuscript, we rewrote this part to better distinguish actual results and our hypotheses.</p>
Excerpt From Revised Manuscript	

-- Ref3.4 – Significance of chromatin remodeling defects --

Reviewer Comment	Papillary RCC with defects in chromatin remodeling genes show a higher mutation burden. This is interesting, but not too surprising as it is the case in other tumor types.
Author Response	To our best knowledge, we are not aware of major systematic studies showing chromatin remodeling (CR) defects are related with higher mutation burden in functionally important DHS regions. Most of the mutation burden studies focus on DNA repair genes. Besides, we showed CR genes mutations are not merely a reflection of high mutation burden but associated directly with mutation landscape change. Our test statistics still stand when the mutation numbers in DHS regions are normalized by the total mutation counts.
Excerpt From Revised Manuscript	

-- Ref3.5 –Methylation analysis--

Reviewer Comment	That methylation influences mutation spectra is interesting and may be pursued, but it needs a more coherent story. Perhaps additional analyses on which mutation pathways are affected and any prognostic role?
Author Response	<p>We thank the reviewer for the suggestions.</p> <ol style="list-style-type: none"> 1. In the revised manuscript, we have added a downstream analysis of methylation-related mutations, emphasizing on the functional consequences of them. 2. During the revision, we realized the alternative splicing event observed in <i>MET</i> in the TCGA study is related to methylation. We showed the novel transcription isoform is due to L1 promoter activation, which is likely due to local hypomethylation. It also reflects global methylation dysfunction. Therefore, the novel <i>MET</i> isoform is associated with methylation cluster 1, which is further away from normal kidney tissues.
Excerpt From Revised	First we validated the TCGA identified methylation cluster 1 showed higher methylation level than cluster 2 in all annotation regions (Figure S2, see Methods), prominently in

Manuscript	<p>CpG Islands (OR of sites being differentially hypermethylated: 1.29, 95%CI: 1.20-1.39, p<0.0001).</p> <p>As expected, C-to-T mutations in CpGs in group 1 showed higher but not statistically significant percentage overlapping with CpG islands compared with group 2 (1.8% versus 1.4%, p=0.14). Therefore, methylation status is the most prominent factor shaping the mutation spectra across patients. We further tried to explore the functional impact of the excessive mutations driven by methylation. C-to-T mutations in CpGs were more likely to be in the coding region (OR=1.54, 95%CI: 1.27-1.85, p<0.0001) and nonsynonymous (OR=1.47, 95%CI: 1.17-1.84, p<0.001). Yet, C-to-T mutations in CpGs did not show functional bias between two methylation groups nor in non-coding regions (Figure SXX).</p> <p>The TCGA study has identified a MET alternative translation isoform as a driver event (3). However, the etiology of this new isoform is unknown. We identified this isoform results from the usage of a cryptic promoter from an L1 element, likely due to a local loss of methylation (REF). This event was reported in several other cancer types (REF). To test its relationship with methylation, we found a closet probe (cg06985664, ~3kb downstream) on the Methylation array show marginally statistically significant (p=0.055, one-side rank-sum test). Additionally, as expected, this event is associated with methylation group 1 (odds ration (OR)= 4.54, 95%CI: 1.07-19.34, p<0.041), indicating genome-wide methylation dysfunction. This association is stronger in type 2 pRCC and it shows a significant association with the C2b cluster (OR= 17.5, 95%CI: 1.72-32.6, p<0.007).</p>
------------	--

-- Ref3.6 – Structural variation analysis --

Reviewer Comment	<p>The structural variations were not explored in great detail. There were 343 SV events but were any recurrent? There were three cases carrying deletions in CDKN2 and 1 case with amplification in MET; otherwise, the structural variations appear as largely a negative result.</p>
Author Response	<p>We understand the concern raised by the referee.</p> <p>First we want to point out that our SV set from sequencing has much finer resolution than the original SNP-array based approach. Therefore, we are able to conduct analyses on breakpoints. Although MET is involved in a lot of amplification events and several samples are genomically unstable, surprisingly we do not find any breakpoint falls into MET and disrupt the gene. This further supports the oncogene role of MET in pRCC.</p> <p>Also, in the revised manuscript, we reanalyze the SVs using a more refined approach. Using high performance cluster, we are able to spend a giant amount of CPU times to realign more than 100 billions of reads for higher quality mapping. We found...</p> <p>We update the manuscript to include the new SV analysis.</p>
Excerpt From Revised Manuscript	<p>First, we recognized that the original BAM files were made by a very old version of BWA that does not support split-read mapping in alignment. Split-read are vital to SV detection. Therefore, we extracted all the reads from the BAMs, paired them and performed remapping. Then we applied LUMPY, a probabilistic SV caller based discordant read pairs and split reads to call the SVs. To evaluate the functional impacts of the somatic SVs, we used SVScore to prioritize and</p>

	evaluate the SVs.
--	-------------------

