

Tags:

Use comma for separation between tags

<ID>	REF 0.0 - title of the comment
<TYPE>	\$\$\$BMR \$\$\$Power \$\$\$Presentation \$\$\$Annotation \$\$\$Network \$\$\$Hierarchy \$\$\$CellLine \$\$\$Stemness \$\$\$Validation \$\$\$NoveltyPos \$\$\$NoveltyNeg \$\$\$Minor \$\$\$Validation \$\$\$Other
<ASSIGN>	@@@XYZ
<PLAN>	&&&AgreeFix - agree and fix &&&DisagreeFix - disagree but we fix, obsequious, and we're safe &&&OOS - out of scope &&&Defer - help me &&&MORE : Go above and beyond the scope of the question and indicates more analyses to be done
<STATUS>	%%%TBC: To Be Continued %%%50DONE: response done (MS+figure to be updated) %%%75DONE: response+calc+figure done (MS to be updated) %%%100DONE: all done. MS+figure+response done

PLEASE NOTE \$\$\$ @@@ &&& %%% are reserved as shown above.
 PLEASE USE ### only for all other tags.

Usage example:

```

<ID>REF 0.0 - Overall comments on the paper
<TYPE>$$$BMR
<ASSIGN>@@@MG,@@@JZ,@@@DL,@@@JL,@@@WM,@@@PDM,@@@Peng,@@
@TG,@@@XK,@@@STL,@@@MTG
  
```

- Style Definition: Heading 1
- Style Definition: Heading 2
- Style Definition: Heading 3
- Style Definition: Heading 4
- Style Definition: Heading 5
- Style Definition: Heading 6
- Style Definition: Title
- Style Definition: Subtitle
- Formatted Table

<PLAN>&&&AgreeFix
<STATUS>%%TBC

Format:

Referee Comment: Courier New
Author Response: Helvetica Neue
Excerpt From Revised Manuscript: Times New Roman

Referee expertise:

Referee #1: cancer genetics, mutational processes
Referee #2: statistical genetics
Referee #3: human genetics
Referee #4: gene expression
Referee #5: cancer genomics

Editor:

<ID>REF 0.1 - Overall comments on the paper

<TYPE>\$\$\$Presentation

<ASSIGN>@@@MG

<PLAN>

<STATUS>%%TBC

[\[JZ2MG: please check the new stuff here. I am also thinking of adding the rewiring here to say cell line is OK, just to highlight the new stuff in this paper\]](#)

Referee Comment	The referees have raised a range of technical concerns on the analyses, including for the background mutation rate, the need to include statistical significance to support many of the claims, and the limitations of this data including cell lines used.
Author Response	<p>We have tried to respond to extensively revise our manuscript in the new version. In summary, we have answered most of these comments. We felt many of them were good suggestions, so we expanded them in large while conserving the manuscript, particularly the suggestions related to</p> <ul style="list-style-type: none">- The overall value of this resource to cancer genomics- Network rewirings from various assays, such RNA-seq, ChIP-seq, and TF knockdowns- Normal-tumor-stem cell comparisons- SVs statistics on networks- Discovery of SUB1 as a potential new oncogene <p>One area that we wish to push back a little on is asking us to compare our calculations to that for driver identification. The point of this paper is not to develop a novel method of driver discovery or to find new cancer drivers. The point is to highlight the use of ENCODE3 data in cancer genomics, particularly related to understanding the overall patterns of mutations, network rewiring, and variant prioritization. Obviously, the ENCODE data will be useful for people developing future driver discovery metrics but we believe that's out of scope for this paper. To respond to previous comments, we have shown how in certain contexts, the ENCODE3 data can help with existing driver discovery measures.</p> <p>Another area we want to mention is the usage of cell lines since some referee preferred tissue data instead of cell lines for cancer. However, as correctly pointed out by referee 4, the genomic and epigenomic heterogeneity in tumor cells, as well as heterogeneity in the tumor microenvironment are significant factors in tumor growth and development. Matching a particular cancer, which is usually quite heterogeneous in nature, to its cell of origin may still be problematic. In our revised manuscript.</p>

Formatted Table

- Deleted: We also
- Deleted: emphasize that although some referees mentioned the limitation of cell line data used here, the
- Deleted: functional genomics data from
- Deleted: of origin is not necessarily a better option
- Deleted: . The
- Formatted: Font:Arial

	<ol style="list-style-type: none"> 1. We tried our best to validate, using external data set, the conclusions we draw from ENCODE call line data and found that our conclusions correlate well with the observations. 2. We clearly pointed it out that ENCODE does not only contain cell line data. For example, 1339 out of 2017 Histone ChIP-Seq experiments we provided for BMR estimation are from primary tissue and we computationally selected the best to use. 3. We added more discussion in the revised manuscript about how technology advances, such as single cell sequencing, can help to provide further insights.
--	---

<ID>REF0.2 – [Regarding context with prior studies](#)

Deleted: Overall comments on the paper

<TYPE>\$\$\$Presentation
 <ASSIGN>@@@MG,@@@JZ
 <PLAN>
 <STATUS>

Referee Comment	The referees also find that the current manuscript provides limited context with prior studies using similar approaches for use of prior ENCODE and Epigenome Roadmap datasets in cancer genomics. They detail the need for clearer presentation in context of prior studies as well comparisons to demonstrate advance.
Author Response	<p>We thank the referees for this comment. We want to note that many of the prior studies have been cited in our initial submission. Some papers, such as Martincorena et al 2017, came out in Nov 2017, two and half months after we submitted our paper in Aug 2017, so it is impossible us to cite in the initial submission. We want to further point that the main focus of the Martincorena et al 2017 paper is not at all about BMR estimation but rather selection patterns in cancer (abstract as below). As suggested, we cited this paper in our revised manuscript and made it clear how our paper is different from this one. However, we feel it is quite unfair for us to make detailed comparisons with it.</p> <p><i>"Universal Patterns of Selection in Cancer and Somatic Tissues: Cancer develops as a result of somatic mutation and clonal selection, but quantitative measures of selection in cancer evolution are lacking. We adapted methods from molecular evolution and applied them to 7,664 tumors across 29 cancer types. Unlike species evolution, positive selection outweighs negative selection during cancer development. On average, <1 coding base substitution/tumor is lost through negative selection, with purifying selection almost absent outside homozygous loss of essential genes. This allows exome-wide enumeration of all driver coding mutations, including outside known cancer genes. On average, tumors carry 4 coding substitutions under positive selection, ranging from <1/tumor in thyroid and</i></p>

Formatted Table

Deleted: In the revised paper, we have clarified the unique aspects of our paper and provided clearer text with previous efforts.

[testicular cancers to >10/tumor in endometrial and colorectal cancers. Half of driver substitutions occur in yet-to-be-discovered cancer genes. With increasing mutation burden, numbers of driver mutations increase, but not linearly. We systematically catalog cancer genes and show that genes vary extensively in what proportion of mutations are drivers versus passengers.](#)

<ID>REF0.3 – [Regarding the advance to the main ENCODE paper](#)

Formatted: Justified
Deleted: Overall comments on

<TYPE>\$\$\$Presentation
<ASSIGN>@@@MG,@@@JZ
<PLAN>&&&DisagreeFix
<STATUS>

Referee Comment	The referees also recommended that the current manuscript does not represent a distinct advance to the main ENCODE manuscript, as it does not report separate new datasets, methods, or clear novel findings. Some referees also recommended that this may be more suitable as Perspective in a specialized journal that further highlights the use on the current ENCODE datasets for cancer genomic studies.
Author Response	<p>We disagree with the reviewers on this point. We want to make it explicit that</p> <p>(1) this paper is to be considered as a "<i>resource</i>" paper, not a novel biology paper</p> <p>(2) the current Encyclopedia <i>package is not meant to be structured like previous packages</i> (i.e. '12 ENCODE). The integrative analysis is meant to be spread over a number of papers and not centered on a single one.</p> <p>(3) note that the ENCODE 3 "data" is not explicitly tied to any paper. Unlike previous roll-outs, ENCODE 3 does not associate particular data sets with specific papers and make use of these data contingent on that paper's publication (as codified in an agreement with NHGRI.)</p> <p>Regarding the novelty of this paper, ENCODEC is unique in its highlighting of a number of ENCODE assays (e.g. replication timing, TF knockdowns, STARR-seq and Hi-C), its deep, integrative annotations combining a wide variety of assays in specific cell types, and its analysis of networks.</p> <p>Note also that while we do NOT feel ENCODEC is a cancer genomics paper, we feel that cancer is the best application to illustrate certain key aspects of ENCODE data and analysis - particularly deep annotations and network changes. We have listed some more details about novelty of this paper as below.</p>

Formatted Table

(1) Networks. These are a core aspect of ENCODE, featured in the '12 roll out. None of the other papers highlight networks in the current package. In ENCODEC, in addition to looking at "universal" ChIP-Seq networks, merged across cell types, we also look at network changes ("rewiring") for specific cell-type comparisons in both proximal and distal networks. We feel that this is best exemplified in oncogenesis.

(2) Deep, integrative annotation – complementary to the Encyclopedia. While the encyclopedia paper considers broad, "universal" annotations across cell-types (currently the centerpiece of ENCODE), it focuses on data common to most cell types (DHS, 2 histone marks and 2 TFs). It does not take advantage of the cell types richer in assays -- the other dimension of ENCODE (diagrammed in ENCODEC's first figure). The ENCODEC paper takes a complementary approach, constructing a more accurate annotation using a large battery of histone marks (>10), next generation assays such as STARR-seq and elements linked by ChIA-PET and Hi-C.

(3) Replication Timing. Although a major feature of ENCODE is replication timing, none of the other papers feature it. Previous work on mutation burden calculation usually selects replication timing data from the HeLa cell line due to the limited data availability. The wealth of the ENCODE replication timing data greatly helps to parametrize somatic mutation rates.

(4) SVs. One unappreciated aspect of ENCODE is that next-generation assays, in addition to characterizing functional elements in the genome, enable one to determine structural variations.

(5) Knockdowns. ENCODE has 222 TF knockout/knockdown experiments, which are not explored systematically in other papers.

Deleted: Excerpt From .

... [1]

Referee #1 (Remarks to the Author):

<ID>REF1.0 – Preamble

<TYPE>\$\$\$Text
<ASSIGN>@@@JZ
<PLAN>&&&AgreeFix
<STATUS>%%%75DONE

We would like to appreciate the referee's feedback. Overall the reviewer mentioned that this is an interesting resource but the novelty of the paper is lacking. We thank the referee for his/her acknowledgement of the potential popularity of our resource for cancer genomics.

Regarding the novelty point, we think differently about the value of our paper. We want to make it clear that [the goal of](#) this paper is to [build a new annotation "resource"](#), not [to discovery](#) novel biology [in cancer](#). We feel that cancer is the best application to illustrate certain key aspects of ENCODE data and analysis - particularly the deep annotations and network changes. We have listed some more details about the resource of this paper as below. Thus, where the referee asks for novelty in cancer gene discovery - we strongly feel that this is out of scope.

Deleted: be considered as

Deleted: a

Formatted: Font:Italic, Underline

Deleted: " paper,

Deleted: paper

Contribution	Subtypes	Data types	ENCODE experiments
Processed raw signal tracks	Histone modification	Signal matrix in TSV format	2015 Histone ChIP-seq
	DNase I hypersensitive site (DHS)	Signal matrix in TSV format	564 DNase-seq
	Replication timing (RT)	Signal matrix in TSV format	51 Repli-seq and Repli-ChIP
	TF hotspots	Signal track in bigWig format	1863 TF ChIP-seq
Processed quantification matrix	Gene expression quantification	FPKM matrix in TSV format	329 RNA-seq
	TF/RBP knockdowns and knockouts	FPKM matrix in TSV format	661 RNAi KD + CRISPR-based KO
Integrative annotation	Enhancer	Annotation in BED format	2015 Histone ChIP-seq 564 DNase-seq STARR-seq
	Enhancer-gene linkage	Annotation in BED format	2015 Histone ChIP-seq 329 RNA-seq

Deleted: -

Formatted Table

	Extended gene	Annotation in BED format	1863 TF ChIP-seq 167 eCLIP Enhancer-gene linkage
SV and SNV callsets	Cancer cell lines	Variants in VCF format	WGS BioNano Hi-C Repli-seq
Network	RBP proximal network	Network in TSV format	167 eCLIP
	Universal TF-gene proximal network	Network in TSV format	1863 TF ChIP-seq
	Tissue-specific TF-gene proximal network	Network in TSV format	1863 TF ChIP-seq
	Tissue-specific imputed TF-gene proximal network	Network in TSV format	564 DNase-seq
	TF-enhancer-gene network level 1-3	Network in TSV format	2015 Histone ChIP-seq 564 DNase-seq

Specifically for the BMR estimation part, the reviewer mentioned that there had been many existing references focusing on applications like cancer driver detection. First, we thank the referee for pointing out to a lot of related references and we did cite many of them in our initial submission. However, some of the references were either published after our initial submission (such as Marticorena et al. 2017) or with a different focus other than BMR estimation (more details in the following table). We updated our reference as suggested but we do not feel it is fair to make a direct comparison for papers with different focuses (JZ2MG: I feel this sentence is too strong). Second, we want to emphasize that the main goal of our paper is not to make a novel driver discovery paper but to illustrate that the richness of the ENCODE data can noticeably help the accuracy of BMR estimation, as we have clearly shown in Fig. 2.

Deleted: . On the reference side, we have listed

Deleted: the papers as the referee suggested and compared

Deleted: with our approach. We have acknowledged the efforts of many of these references.

Deleted: was out after our initial submission so we did not have a chance to add them. In the revised version we have further expanded our reference list for some the publications

Deleted: date. We

Deleted: actually

Deleted: many of the methods used in these papers. With a larger pool of covariate selection, the estimation

Deleted: can be significantly improved.

Formatted: Font color: Black

Reference	Initial	Revised
Lawrence et al, 2013	Cited	Cited
Weinhold et al, 2014	Cited	Cited
Araya et al, 2015	No	Cited
Polak et al (2015)	Cited	cited
Martincorena et al (2017)	No (out after our submission)	Cited
Imielinski (2017)	No	Yes
Tomokova et al. (2017)	No	Yes
huster-Böckler and Lehner (2012)	Yes	Yes
Frigola et al. (2017)	No	Yes
Sabarinathan et al. (2016)	No	Yes
Morganella et al. (2016)	No	Yes
Supek and Lehner (2015)	No	Yes

Deleted:

Reference	Initial	Revised	Main point	Comments
Lawrence et al, 2013	Cited	Cited	Introduce replication timing and gene expression as covariates for BMR correction	Replication timing in one cell type
Weinhold et al, 2014	Cited	Cited	One of the first WGS driver detection over large scale cohorts.	Local and global binomial model
Araya et al, 2015	No	Cited	Sub-gene resolution burden analysis on regulatory elements	Fixed annotation on all cancer types
Polak et al (2015)	Cited	cited	Use epigenetic features to predict cell of origin from mutation patterns	Use SVM for cell of origin prediction, not specifically for BMR
Martincorena et al (2017)	No (out after our submission)	Cited	Use 169 epigenetic features to predict gene level BMR	No replication timing data is used
Imielinski (2017)	No	Yes	Use ENCODE A549 Histone and DHS signal for BMR correction	Limited data type used from ENCODE
Tomokova et al. (2017)	No	Yes	8 features (5 from ENCODE) for BMR prediction and mutation/indel hotspot discovery	Expand covariate options from ENCODE data
huster-Böckler and Lehner (2012)	Yes	Yes	Relationship of genomic features with somatic and germline mutation profiles	NOT specifically for BMR
Frigola et al. (2017)	No	Yes	Reduced mutation rate in exons due to differential mismatch repair	NOT specifically for BMR
Sabarinathan et al. (2016)	No	Yes	Nucleotide excision repair is impaired by binding of transcription factors to DNA	NOT specifically for BMR
Morganella et al. (2016)	No	Yes	Different mutation exhibit distinct relationships with genomic features	NOT specifically for BMR
Supek and Lehner (2015)	No	Yes	Differential DNA mismatch repair underlies mutation rate variation across the human genome.	NOT specifically for BMR

<ID>REF1.1 – [Positive comments](#) on the resource releases

Deleted: Comments

<TYPE>\$\$\$NoveltyPos
 <ASSIGN>
 <PLAN>&&&AgreeFix
 <STATUS>%%%75DONE

Referee Comment	This manuscript describes how the ENCODE project data could be utilized to derive insights for cancer genome analysis. It has several examples to illustrate this point, e.g., how to
-----------------	---

Formatted Table

	better estimate background mutation rate in a cancer genome, how to modify gene annotation for finding mutation-enriched regions (e.g., by bundling enhancer regions to target genes using Hi-C/ChIA-PET), and describing the changes in regulatory networks in cancer. Obviously, the ENCODE project involves a great deal of planning and a lot of experimental work by many groups, and the overall aim of re-highlighting the ENCODE as a resource to cancer research seems worthwhile in general, perhaps even in a high-profile journal.
Author Response	We thank the referee for the positive feedback.

<ID>REF1.2 – BMR: comparison with existing literature

<TYPE>\$\$\$BMR,\$\$\$Text

<ASSIGN>@@@JZ,@@@WM,@@@PDM

<PLAN>&&&OOS

<STATUS>%%%75DONE

Referee Comment	Just to take the first application as an example, the problem of estimating background somatic mutation rate accurately in order to better identify cancer drivers has been studied extensively in the literature. One paper, "Mutational heterogeneity in cancer and the search for new cancer-associated genes" (Nature 2013), is cited in the current manuscript, but there are many others. For instance, Weinhold et al, 2014 (Genome-wide analysis of noncoding regulatory mutations in cancer, Nat Genetics), Araya et al, 2015 (Identification of significantly mutated regions across cancer types highlights a rich landscape of functional molecular alterations, Nat Genetics), and similar non-coding mutation identification papers all include steps to account for epigenetic features in their background rate calculation.
Author Response	We thank the reviewer for identifying these references. We <u>did</u> recognize that genomic features <u>were</u> used to estimate BMR and improve driver mutation detection. Our aim here was <u>not to</u> claim a better BMR estimation model nor <u>to propose</u> a novel discovery that "matched" features performs better. We made it

Formatted Table

Deleted: have been previously been

Deleted: neither

Deleted: claim

	<p>more apparent in our revised manuscript that our purpose is to showcase how ENCODE data can help BMR estimation in many models.</p> <p>With the wealth data available through ENCODE data, we had a much larger pool of features to choose from to potentially improve BMR estimation. There are thousands of histones modification marks that are released into a ready to use format (see details in the table below).</p> <p>Also, we have provided other data types, such as replication timing, that has been proven to affect BMR but has not been widely by others. We believe that such data, when released into a ready to format, can help BMR estimation through many existing models.</p>
Excerpt From Revised Manuscript	

Deleted: clearer

Formatted: Justified

Deleted: In addition

Deleted: have

Deleted: be

Deleted: have

Cell Type	# t
tissue	81
primary-cell	52
cell-line	33
in-vitro-differentiated-cells	17
stem-cell	11
induced-pluripotent-stem-cell-line	46

<ID>REF1.3 – BMR: [Match](#)

<TYPE>\$\$\$BMR,\$\$\$Text
 <ASSIGN>@@@JZ,@@@WM
 <PLAN>&&&DisagreeFix
 <STATUS>%%%50DONE

Deleted:

Deleted: lacking in novelty in the conclusion

Deleted: 75DONE

Formatted Table

Deleted: thank

Deleted: referee for pointing out

Deleted: Polak 2015 paper. This is an important reference to relate various genomic features to cancer mutational landscape, and we did cite

Deleted: in our initial submission. Excerpt From - ... [2]

Deleted: discoveries in that paper,

Deleted: help BMR estimation. We also want to emphasize that two points here. Excerpt From - ... [3]

Formatted: Font:Helvetica Neue, No underline

Deleted: does not exclude that other "non-matched" features from being useful to improve the BMR prediction accuracy. Actually some of the recent papers, such Martincorena et al

Moved down [1]: (2017), also used the top 20 PCs of 169 histone features in their model. On this point, we uniformly processed thousands of features in a ready-to-use format. Many of them are not mentioned in other literature, such as replication time from 51 tissue/cell lines. They have proven useful but are less frequently matched probably due to the lack of data incorporated into previous BMR models

Formatted: Font:Helvetica Neue

Deleted: Excerpt From - ... [4]

Referee Comment	<p>Most large-scale cancer genome sequencing papers also have models at various levels sophistication, most of them including the issue of proper tissue-type matching. "matched" cell lines are better than unmatched or addition of more epigenetic features results in some improvement is almost trivial at this point. Which marks contribute to this is also not new.</p>
Author Response	<p>We think differently about the purpose of the BMR section. Please note that the goal of this paper is to build a new annotation "resource" using the richness of ENCODE data. Here, we are not trying to reproduce the claims on how epigenomic features affect BMR but rather to show how the richness of ENCODE data can make improved BMR estimations.</p>

<ID>REF1.4 – BMR: Tissues vs. Cell lines

<TYPE>\$\$\$BMR,\$\$\$Calc

<ASSIGN>@@@JZ,@@@JL

<PLAN>&&&DisagreeFix,&&&More

<STATUS>%%%50DONE

Referee Comment	<p>Importantly, Polak et al, 2015 (Cell-of-origin chromatin organization shapes the mutational landscape of cancer, Nature) in fact show that cell-of-origin chromatin features are much stronger determinants of cancer mutations profiles than chromatin feature of matched cancer cell lines, and that cell type origin can be predicted from the mutational profile.</p> <p>Stepping back, it is not obvious to me that using the ENCODE cell lines, despite the availability of more epigenetic data, is the best approach to calculating the background rate in the first place—they briefly mention that using cell lines (rather than tissues) can be problematic, but do not explore this further. If this were a regular research paper, the authors would have to shown how the proposed approach is different and how it is better than methods already available.</p>
Author Response	<p>We thank the referee for pointing out the comparison of cell line vs. tissues and we feel this is a good suggestion. In our revised manuscript, we further investigated it in detail by extending our analysis to many new data types, such as RNA-seq and distal/proximal TF ChIP-Seq data. We think slightly differently with the referee on value of cell line data. Several points we want to emphasize are</p> <p>- On a large scale (up to mbp)</p> <ul style="list-style-type: none"> • First, the Polak 2015 paper did not perform large-scale comparison across various cancer cell lines. As seen from Except 1 below, cell line data provides comparable, sometimes even better, correlation with mutation counts. We have added a new section in the supplementary file to discuss this. • As compared to cell line data, there are way less functional characterization data in tissues. For example, there are no prostate tissue data from the REMC. We have updated supplementary table 1 for a comparison of data richness in ENCODE3.

Deleted: 75DONE

Deleted: - ... [5]

Formatted Table

Deleted: tissue. We

Deleted: this comparison and extended this point more

Deleted: the

Deleted: this point.

Formatted: Outline numbered + Level: 1 + Numbering Style: Bullet + Aligned at: 0.25" + Indent at: 0.5"

Deleted: the following figure

Deleted: (such as

Deleted:).

	<ul style="list-style-type: none"> • ENCODE is not just about cell lines, and there are many ENCODE tissue data for histones (339 cell line vs 818 tissue, details see excerpt 2 below). We have added a supplementary table on this point. • The fact that "matched" cell type features performs better in predicting BMR does not exclude other "non-matched" features from being useful to improve the BMR prediction accuracy. Actually some of the recent papers, such as Martincorena et al. (2017), also used the top 20 PCs of 169 histone features in their model. On this point, we uniformly processed thousands of features in a ready-to-use format. Many of them are not mentioned in other literature, such as replication time from 51 tissue/cell lines. They have proven useful but are less frequently matched probably due to the lack of data incorporated into previous BMR models. We believe that this is quite useful for cancer genomics. <p>- On a small scale (less than kbp) Features, like expression levels and TF binding events, have been used widely to affect somatic mutation rates. As suggested by the referee, we systematically investigated the RNA-seq and TF ChIP-Seq data and found that many of the cancer transcriptome/TF binding landscape are quite similar to each other, as compared to the initial of primary cells. This has also been mentioned by previous reports, such as Lotem et al. 2005 and Hoadley et al. 2014. The fact that cancer cells lose diversity and showed a distinct pattern from the primary cells highlights the values of cell line data. We have added this result into the main figure and supplementary files.</p>
<p>Excerpt 1 From Revised Supplementary file</p>	<p>1. Comparison of mutation rate vs features in tissue/cell lines. We provided the pearson correlation of the breast cancer mutations count per Mbp vs. various histone modification features in tissue and cell line. Cell line data provides comparable (and sometimes even better) correlation with mutation counts.</p>

Deleted:).
Formatted: Font:Bold

Moved (insertion) [1]

Deleted: -

Deleted: Manuscript

	<p style="text-align: center;">BRCA var counts/mbp vs Histone Sig/mbp</p> <table border="1"> <caption>Approximate Pearson Correlation values from the chart</caption> <thead> <tr> <th>Histone Sig</th> <th>MCF-7</th> <th>HMEC</th> </tr> </thead> <tbody> <tr> <td>H3K27me3</td> <td>0.35</td> <td>0.32</td> </tr> <tr> <td>H3K36me3</td> <td>-0.35</td> <td>-0.45</td> </tr> <tr> <td>H3K4me1</td> <td>-0.35</td> <td>-0.40</td> </tr> <tr> <td>H3K4me3</td> <td>-0.30</td> <td>-0.35</td> </tr> <tr> <td>H3K9me3</td> <td>0.35</td> <td>0.32</td> </tr> </tbody> </table>	Histone Sig	MCF-7	HMEC	H3K27me3	0.35	0.32	H3K36me3	-0.35	-0.45	H3K4me1	-0.35	-0.40	H3K4me3	-0.30	-0.35	H3K9me3	0.35	0.32
Histone Sig	MCF-7	HMEC																	
H3K27me3	0.35	0.32																	
H3K36me3	-0.35	-0.45																	
H3K4me1	-0.35	-0.40																	
H3K4me3	-0.30	-0.35																	
H3K9me3	0.35	0.32																	
<p>Excerpt 2 From Revised Supplementary file</p>	<p>2. Summary of ENCODE histone ChIP-seq data</p> <table border="1"> <thead> <tr> <th>Cell Type</th> <th># histone marks</th> </tr> </thead> <tbody> <tr> <td>tissue</td> <td>818</td> </tr> <tr> <td>primary-cell</td> <td>521</td> </tr> <tr> <td>cell-line</td> <td>339</td> </tr> <tr> <td>in-vitro-differentiated-cells</td> <td>179</td> </tr> <tr> <td>stem-cell</td> <td>114</td> </tr> <tr> <td>induced-pluripotent-stem-cell-line</td> <td>46</td> </tr> </tbody> </table>	Cell Type	# histone marks	tissue	818	primary-cell	521	cell-line	339	in-vitro-differentiated-cells	179	stem-cell	114	induced-pluripotent-stem-cell-line	46				
Cell Type	# histone marks																		
tissue	818																		
primary-cell	521																		
cell-line	339																		
in-vitro-differentiated-cells	179																		
stem-cell	114																		
induced-pluripotent-stem-cell-line	46																		
<p>Excerpt 3 From Revised Supplementary file</p>	<p>3. We performed RCA/PCA analysis on RNA-Seq, shRNA RNA-Seq, and TF ChIP-seq data and found that cancer cells demonstrate a consistent pattern to be more similar to stem cells, as compared to their primary cells of origin.</p>																		

BRCA var counts/mbp vs


Histone Sig	MCF-7	HMEC
H3K27me3	0.35	0.32
H3K36me3	-0.35	-0.45
H3K4me1	-0.35	-0.40

Deleted:

Deleted:

Cell Type	#
tissue	8
primary-cell	5
cell-line	3
in-vitro-differentiated-cells	1
stem-cell	1
induced-pluripotent-stem-cell-line	4
... [6]	

Moved (insertion) [2]



REF1.5 – Difference between ENCODEC and Prev. prioritization methods

<TYPE>\$\$\$BMR,\$\$\$Text
 <ASSIGN>@@@JZ
 <PLAN>&&DisagreeFix
 <STATUS>%%%75DONE

Referee Comment	The rest of the sections (and their corresponding supplement sections) are variable in significance and quality. That ENCODE data helps in prioritization of non-coding variants has been well demonstrated already (including by some of the authors on this paper), and so the value of the described analysis less clear.
Author Response	The referee pointed out that we and others have tried to prioritize non-coding elements before. This is definitely true and we are not claiming to be the first. However, we believe that the method that we used here is new and novel. The important aspect is that it takes advantage of many new ENCODE data and integrates over many different aspects. Detailed changes please see the Excerpt blow.
Excerpt From Revised Manuscript	In particular, it takes into account the STARR-seq data, the connections from Hi-C, the better background mutation rates, and the network rewiring data, which is only possible in the context of the highly integrated and their data available on certain cell lines. We are showing this as an example of the best we can do with this level of integration. The fact that we coupled this with quite successful validation that we believe points to the great value of the integrated incurred data.

Deleted: 100DONE

Formatted Table

Deleted: -

... [7]

Deleted: In particular, it takes into account the STARR-seq data, the connections from Hi-C, the better background mutation rates, and the network wiring data, which is only possible in the context of the highly integrated and their data available on certain cell lines.

Moved down [3]: We are showing this as an example of the best we can do with this level of integration. The fact that we coupled this with quite successful validation that we believe points to the great value of the integrated incurred data.

Formatted: Font:Times New Roman

Moved (insertion) [3]

Formatted: Font:Times New Roman

<ID>REF1.6 – Novelty and presentation of the paper

<TYPE>\$\$\$Presentation,\$\$\$NoveltyPos,\$\$\$NoveltyNeg,\$\$\$Text

<ASSIGN>@@@JZ

<PLAN>&&&AgreeFix

<STATUS>%%%50DONE

[\[JZ2MG: should we only preserve the starr-seq comments here and remove the lack of novelty here? The novelty issues are all in the preamble?\]](#)

Referee Comment	Some newer assays such as STARR-seq are helpful, obviously, in better predicting enhancers, but, again, while the analysis done serves as illustrations how ENCODE data can be used, the supplement does not seem to give a convincing evidence of how the results found are novel.
Author Response	<p>We thank the referee for praising the novel assays, such as STARR-seq, and we have in fact tried to illustrate the value of novel assays such as STARR-Seq. We have modified both the main manuscript and the supplement to further highlight this.</p> <p>As for the enhancer part, with the ensemble method, for example, we can get more accurate annotation and pin-point to sequences where transcription factors would actually bind to. To estimate the false positive rate would not be very practical at this stage as there is no gold-standard experiment that could assert an predicted enhancer is definitely negative. Here we took the FANTOM enhancer data set and assess the overlap percentage of our enhancer annotation in each ensemble step. We show that each ensemble step indeed increases the percentage of overlap between our annotation and the FANTOM enhancer set. The overlap percentage for our annotation is much higher than that of the Roadmap annotation, and is also higher than the main encyclopedia enhancer annotation annotation (ccRE).</p>
Excerpt From Revised Manuscript	We have performed QC of different types of enhancers in details in K562 and GM12878 as an example to show the power of integrating various types of assays.



Deleted: DONE

Deleted: JZ2MTG: would u pls update the figure? The legend is too small to see and would you please change it to a barplot? .

Formatted Table

Deleted: new

Deleted: assays



Referee #2 (Remarks to the Author):

<ID>REF2.0 – Preamble

<TYPE>\$\$\$Text
 <ASSIGN>@@@MG,@@@JZ
 <PLAN>&&&AgreeFix
 <STATUS>%%75DONE

We would like to appreciate the referee's feedback, especially about the positive comments on the value of our resource, the extended gene, and the network rewirings. Regarding the novelty point, [we want to emphasize that](#) this paper is unique in its highlighting of a number of ENCODE assays (e.g., replication timing, TF/RBP knockdowns, STARR-seq, ChIA-PET, and Hi-C), its deep, integrative annotations combining a wide variety of assays in specific cell types, and its analysis of networks. Note also that while we do NOT feel this is a cancer genomics paper, we feel that cancer is the best application to illustrate certain key aspects of ENCODE data and analysis - particularly deep annotations and network changes. We have listed some more details about the novelty of this paper as below.

Deleted: Regarding the novelty of our work,
 Deleted: .

Contribution	Subtypes	Data types	ENCODE experiments
Processed raw signal tracks	Histone modification	Signal matrix in TSV format	2015 Histone ChIP-seq
	DNase I hypersensitive site (DHS)	Signal matrix in TSV format	564 DNase-seq

Formatted Table

	Replication timing (RT)	Signal matrix in TSV format	135 Repli-seq and Repli-ChIP
	TF hotspots	Signal track in bigWig format	1863 TF ChIP-seq
Processed quantification matrix	Gene expression quantification	FPKM matrix in TSV format	329 RNA-seq
	TF/RBP knockdowns and knockouts	FPKM matrix in TSV format	661 RNAi KD + CRISPR-based KO
Integrative annotation	Enhancer	Annotation in BED format	2015 Histone ChIP-seq 564 DNase-seq STARR-seq
	Enhancer-gene linkage	Annotation in BED format	2015 Histone ChIP-seq 329 RNA-seq
	Extended gene	Annotation in BED format	1863 TF ChIP-seq 167 eCLIP Enhancer-gene linkage
SV and SNV callsets	Cancer cell lines	Variants in VCF format	WGS BioNano Hi-C Repli-seq
Network	RBP proximal network	Network in TSV format	167 eCLIP
	Universal TF-gene proximal network	Network in TSV format	1863 TF ChIP-seq
	Tissue-specific TF-gene proximal network	Network in TSV format	1863 TF ChIP-seq
	Tissue-specific imputed TF-gene proximal network	Network in TSV format	564 DNase-seq
	TF-enhancer-gene network level 1-3	Network in TSV format	2015 Histone ChIP-seq 564 DNase-seq

<ID>REF2.1 – Comment on utility of the resource

<TYPE>\$\$\$NoveltyPos

<ASSIGN>

<PLAN>&&&AgreeFix

<STATUS>%%100DONE

Referee Comment	However, there is a possibility that the resource would be very popular among cancer genomics researchers. Also, results on extended genes and rewiring are of interest.
Author Response	We thank the referee for the positive comment.

Formatted Table

<ID>REF2.2 – Comparison of negative binomial to other methods

<TYPE>\$\$\$BMR,\$\$\$Text,\$\$\$Calc

<ASSIGN>@@@JZ

<PLAN>&&&OOS

<STATUS>%%%75DONE

Deleted: 100DONE

Referee Comment	1) The negative binomial regression (Gamma-Poisson mixture model) was introduced in Nik-Zainal et al. Nature 2016 and Marticorena et al., Cell 2017. Why was not this available method applied, and what is the benefit for the procedure used by the authors?
Author Response	<p>The referee is pointing out that negative binomial regression has been used before. This is a standard statistical technique that has been used in many contexts. Please note that the Marticorena et al. paper came out in Nov 2017, which was almost three months after our initial submission and it is more about positive selection instead of BMR estimation. The fact that it also used negative binomial regression bolsters the underlying technical validity of our argument. While we admit it does slightly undercut a claim of novelty in this regard, that is not central to our work.</p> <p>Also, ENCODE3 provides noticeably more covariate data, which is uniformly processed and less explored in the references mentioned by the referees. Some features, such as replication timing, that is well-known confounders but was not included in the Marticorena et al. paper. We are not aiming to make a new method for predicting background mutation rate, but rather to use a robust regression method that really takes into account the very large amount of data and is able to leverage that to more successfully predict background mutation. Therefore, we did not directly use their approach.</p>

Formatted Table

Deleted: The fact that the recent Marticorena et al 2017 paper uses this, we think only

Deleted: that's

Deleted: There is new data type

Deleted: ,


Deleted: those papers. Our

Deleted: is


Deleted: . [8]

<ID>REF2.3 – Questions about the Goodness of fit of the Gamma-Poisson Model

<TYPE>\$\$\$BMR,\$\$\$Calc
 <ASSIGN>@@@JZ
 <PLAN>&&&AgreeFix,&&&OOS
 <STATUS>%%100DONE


Referee Comment	Also, does Gamma-Poisson model fits data for most cancers well or is it just an approximation? One can use non-conjugate priors but this is probably beyond the scope of this work.
Author Response	We thank the referee for mentioning the goodness of fit of the Gamma-Poisson model. As suggested, we provided more figures in our supplementary file to investigate this. For most of the cancer types, the fitting of Gamma-Poisson is pretty good (as seen in the figures below). Also, we point out the fact that it has been used in other literature provides further technical support for this using. However, we agree that it is interesting to investigate other non-conjugate priors. As the referee mentioned, this is out of scope, but we have made a mention of this in the text.
Excerpt From Revised Supplementary file	

Formatted Table



Deleted:
Deleted: Manuscript

... [9]



<ID>REF2.4 – Was the Poisson Model used for low mutation cancers

<TYPE>\$\$\$BMR,\$\$\$Text,\$\$\$Cale

<ASSIGN>@@@JZ,@@@JL

<PLAN>&&&AgreeFix

<STATUS>%%%75DONE

Deleted: 100DONE

Referee Comment	2) It seems that the Poisson model was not rejected for cancers with very low mutation counts (liquid tumors). Is this a power issue rather than the property of the mutation process?
-----------------	--

Formatted Table

Referee Comment	3) The approach with principal components used for the BMR estimation does not seem to work well. Starting with the second PC most components have roughly the same prediction power. One possibility is that higher principle components do not capture the additional signal and reflect noise in the data, and the correlation with mutation rate is due to an overfit of the NB regression (it is unclear whether it was analyzed with cross-validation). Another possibility is that the signal is spread over many components. In the latter case, this is not an optimal method choice.
Author Response	<p>We thank the referee for pointing out the limited contribution from the higher order principal components. In fact, we wanted to bring out this point, and we <u>do not</u> see this as efficient either. The point of our approach is not to say that a few top components or a few features can predict a mutation rate accurately. Actually we want to show the opposite that the wealth of the ENCODE data is useful and that with additional data types, one gets a small but measurable continued improvement. We use principal components essentially as a way of doing a principled unbiased feature selection, but we realized that actually <u>did not</u> get across very clearly, so we have replotted this figure and now simply show how one gets <u>a</u> steady increase in predictions forms by just adding features one at a time.</p> <p>We hope this gets the point across. The aim here is <u>not to</u> highlight a complicated mathematical method but just simply to get across the idea that the <u>extensive</u> ENCODE data provides a valuable resource for predicting BMR and we appreciated the referee helping us achieve clarity on this point. We put the main text figures into the supplementary files and made for the main.</p>
Excerpt From Revised Manuscript	<ol style="list-style-type: none">1. At 1mb bin resolution, we compared the performance of models using random features vs. computationally selecting best features sequential (forward selection). It has shown that by adding features appropriately from ENCODE3, we can noticeably improve the performance of BMR accuracy.

Formatted Table

Deleted: actually

Deleted: don't


Deleted: didn't

Deleted: to



Deleted: very large

Deleted: -


Formatted: Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"



2. To avoid overfitting problem, we performed 5 fold cross validation using the selected model for each cancer type and listed the performance as below.



Deleted:



Deleted:

<ID>REF2.6 – Comments on the power analysis and compact annotations

<TYPE>\$\$\$Power,\$\$\$Calc

<ASSIGN>@@@JZ

<PLAN>&&&AgreeFix

<STATUS>%%%75DONE

[\[JZ2JZ:wait for the GWAS to be added here\]](#)

Referee Comment	<p>4) I do not agree with the power analysis presented to support the idea of compact annotations. I understand that this is a toy analysis neglecting specific properties of mutation rate known for regulatory regions and also sequence context dependence of mutation rate. The larger issue is that the analysis assumes that ALL functional sites are within the compact annotation. In that case, power indeed would decrease with length. <u>However, in case some of the functional sites are outside the compact annotation power would not decrease and is even likely to increase with the inclusion of additional sequence.</u> Is there a justification for all functional sites to reside within compact annotations? Can this issue be explored? Some statistical tests incorporate weighting schemes.</p>
Author Response	<p>The referee is indeed correct and we expanded our power calculation in our revised manuscript. In our initial submission, the assumption is that we were trimming off the nonfunctional sites while preserving the functional ones. Two examples can explain the motivation of this assumption, (see details in excerpt 1 below).</p> <p>▲ Following the reviewer's suggestions, in our revised manuscript we show in a formal power analysis that the most important contribution to power comes from including additional functional sites, which is of course by the extended gene concept and then secondarily, from removing non-functional sites, but to a lesser extent. The assumption in our compacting annotations is that we can accurately distinguish the more important functional nucleotides from the less important ones through the guidance of many functional characterization assays.</p> <p>Admittedly, we are making assumptions and the referee is completely correct in pointing this out. We have tried to be more precise in the text that we are assuming that the large number of ENCODE assays, when integrated, allow us to more</p>

Formatted Table

Deleted: .

Moved down [5]: -

... [11]

	<p>directly get the functional nucleotides, but this, of course, is an assumption. It is hard to tell to what degree one can succeed in finding the current events in cancer. It is hard to back this up with the gold standard, but we think that some of the points are self evidently obvious. We have tried to make this clear in text and thank the referee for pointing this out.</p>
<p>Excerpt 1 From Revised Supplementary file</p>	<p>Two examples can explain the motivation of this assumption.</p> <p>1) Enhancers: Traditionally, enhancers were called as a 1kb peak regions, which admittedly introduced a lot of obviously nonfunctional sites. We believe we can get functional region more accurately by trimming the enhancers down using the exact shapes of many histone marks and further integration with STARR-seq and Hi-C data.</p> <p>2) TFBS hotspots around the promoter region of WDR74. Instead of testing the conventional up to 2.5K promoter region, we can trim the test set to a core set of the promoter region where many TFs bind, which perfectly correlates with the mutation hotspots (red block) for this well-known driver site (blue line for pancreatic and green line for liver cancer).</p>
<p>Excerpt 1 From Revised Manuscript</p>	<p>GWAS for power analysis</p>

Moved (insertion) [5]

Formatted Table

Deleted:

REF2.7 – Q-Q plots

<TYPE>\$\$\$BMR,\$\$\$Calc

<ASSIGN>@@@JZ

<PLAN>&&&Defer

<STATUS>%%TBC

####Thinking

[JZ2MG: not finished yet for this part]

Referee Comment	5) Some of the QQ-plots in supplementary figures look problematic. Also, for some tumors with low count statistics QQ-plots are expected to always be deflated, so the interpretation of QQ-plots may be non-trivial.
Author Response	This is a good point. We've done XXX & YYY now But we wish to make clear that the point of this paper is not driver detection Our goal is BMR We show QQ w diff detection We actually show QQ plots with drivers Take some else's driver detection method, use our BMR model, show that it works better
Excerpt From Revised Manuscript	

Formatted Table

<ID>REF2.8 – Value of the extended gene

<TYPE>\$\$\$NoveltyPos

<ASSIGN>

<PLAN>&&&AgreeFix,&&&MORE

<STATUS>%%%50DONE

Deleted: 75DONE

Deleted: [JZ2JL: please add your figure here]

Referee Comment	6) The idea of extended genes and the use of multiple information sources to construct them is a strength of the paper.
Author Response	We thank the reviewer for the positive remarks. We further highlighted this part in our revised manuscript and added several new sections to highlight the value of extended genes, such as 1. We extensively expanded our power analysis part to include more extended gene analysis (as we pointed up in the response to <ID>REF2.6 – Comments on the power analysis and compact annotations)

Formatted Table

	2. We showed that by using the extended gene, we can better stratify the gene expressions.
Excerpt From Revised Manuscript	JL figure to be added here on Monday More to added from the GWAS side

Deleted: -

<ID>REF2.9 – BMR effect on local tri-nucleotide context

Deleted: 10

<TYPE>\$\$\$BMR,\$\$\$Text

<ASSIGN>@@@JZ

<PLAN>^^^AgreeFix


<STATUS>%%%75DONE

Deleted: 100DONE

Referee Comment	However, it is unclear whether the analysis takes into account complexities of the mutation model in regulatory regions. The influence of tri- or even penta-nucleotide context can be significant.
Author Response	In the main figure, we did not show how local context effect may affect BMR in order to highlight the effect of accumulating features. However, in the supplementary file where we described our method, we separate the 3mers to run negative binomial regression. We showed that in Supplementary figure xxx that local context effect is huge - usually up to several order of effect on BMR. (Please see details in the following excerpt).
Excerpt From Original Supplementary file	Consistent with previous literature, we observed large mutational heterogeneity over the genome for all 3-mers in all cancer types. As seen in Figure S 2-2 , the mutation rate changes significantly over different regions of the genome (large region of each violin bar) and over different local contexts.

Formatted Table

Deleted: . We made this point more clear in our revised manuscript.



<ID>REF2.10 – Confounding factors

Deleted: 11

<TYPE>\$\$\$Other

Deleted: BMR

<ASSIGN>@@@JZ

<PLAN>&&&AgreeFix

<STATUS>%%%75DONE

Deleted: 100DONE

Referee Comment	Next, TF binding and nucleosome occupancy is known to interfere with the activity of DNA repair system.
Author Response	We thank the referee to bring out this important point. Actually many of the current background mutation rate estimation method assumes a constant rate in a fairly large region, such as a within a gene (including the long introns in between) or up to Mbp fixed bins. In such large scale, it is difficult to incorporate such as TF binding, nucleosome occupancy, histone modification (which changes sharply in less kbps). Hopefully, with accumulating cancer patient data in the future could help to build up site specific background models to investigate more about such effects. We added this point in our discussion section.

Formatted Table

Excerpt From Revised Manuscript	Hower, most of the current BMR models are focused on larger scale mutation rate variations by integrating many features at 50 kb to 1 Mb resolution while ignoring small scale perturbations introduced by TF binding and nucleosome occupancy. Improvement of such finer scale features in the future could further improve BMR estimation.
---------------------------------	--

<ID>REF2.11 – Power analysis of extended genes

Deleted: 12

<TYPE>\$\$\$Power,\$\$\$Calc
 <ASSIGN>@@@JZ
 <PLAN>&&AgreeFix
 <STATUS>%%75DONE

Referee Comment	It would be great to see a formal analysis about how extended genes increase power of cancer driver discovery.
Author Response	We thank the referee for this comment and encouraging us to do a formal analysis. We have expanded our power analysis in the revised manuscript.
Excerpt From Revised Manuscript	We showed in a formal power analysis that the most important contribution to power comes from including additional functional sites, which is of course by the extended gene concept and then secondarily, from removing non-functional sites, but to a lesser extent. The assumption in our compacting annotations is that we can accurately distinguish the more important functional nucleotides from the less important ones through the guidance of many functional characterization assays. Admittedly, we are making assumptions and the referee is completely correct in pointing this out. We have tried to be more precise in the text that we are assuming that the large number of ENCODE assays, when integrated, allow us to more directly get the functional nucleotides, but this, of course, is an assumption. It is hard to tell to what degree one can succeed in finding the current events in cancer. It is hard to back this up with the gold standard, but we think that some of the points are self evidently obvious. We have tried to make this clear in text and thank the referee for pointing this out.

Formatted Table

Deleted: - ... [12]

<ID>REF2.12 – Minor comment on burden test

Deleted: 13

<TYPE>\$\$\$Minor,\$\$\$Presentation,\$\$\$Text
 <ASSIGN>@@@JZ
 <PLAN>&&AgreeFix

<STATUS>%%75DONE

Referee Comment	1) I would not use the term "burden test". This usage is slightly confusing because this term is commonly used in human genetics where it refers to a case-control test.
Author Response	We thank the referee to point out this. We have changed our terminology in our revised manuscript.

Formatted Table

<ID>REF2.13 – Minor comment on terminology

<TYPE>\$\$\$Minor,\$\$\$Presentation,\$\$\$Text

<ASSIGN>

<PLAN>&&AgreeFix

<STATUS>%%75DONE

Deleted: 14

Referee Comment	2) Similarly, it is unclear what is meant by "deleterious SNVs" as the term is commonly used in human genetics in reference to germline variants under negative selection.
Author Response	We thank the referee to point out this. "Deleterious SNVs" in our manuscript means somatic mutations that disrupts gene regulations. To avoid potential confusion, we changed it in our revised manuscript.

Formatted Table

Referee #3 (Remarks to the Author):

<ID>REF3.0 – Preamble

<TYPE>\$\$\$Text
<ASSIGN>@@@MG,@@@JZ
<PLAN>&&&AgreeFix
<STATUS>%%%75DONE

In relation to the supplement and genomics, the referee points out that it's sometimes hard to see full documentation of our methods in the main part and one has to look at the extensive supplements. We are well aware of this fact. The very large scale of supplement is typical for large genomic paper. We, in fact, have been actively discussing with Nature Publishing and other companions about the supplement with regard to the main text. We have attempted to put important things in the supplement and to structure it very carefully. We admit that maybe this construction is not that intuitive. We are prepared to work very hard to make the structure of the supplement understandable. We've tried to revise it to make these clearer and also to move more appointives into the main text, though we think given the current main text limitations of a typical paper nature and the scale of the results in the data in this paper, it's simply impossible to put everything into the main text. We are preparing to work constructively with the referees and the others to make this clear.

<ID>REF3.1 – Presentation of the paper

<TYPE>\$\$\$Presentation
<ASSIGN>
<PLAN>&&&AgreeFix
<STATUS>%%%TBC

Referee Comment	It is difficult to understand the significant novel findings in this paper (compared to the main ENCODE paper). Perhaps, some of this is due to the data not being presented in a concise and clear manner. For example, I wonder whether the authors can add more details and straightforward directions when citing supplementary information. In the current main manuscript, the authors cited all supplementary information as (see suppl.). It might be hard for the reader to check where the authors refer to in the supplementary information. I think more direction, such as sup Fig1, sup Table 1, or section 7.2S etc, would be very helpful.
-----------------	--

Formatted Table

Author Response	We tried the new way of citing supplementary info.
Excerpt From Revised Manuscript	

<ID>REF3.2 – Benefits of using multiple cancer types in BMR

<TYPE>\$\$\$BMR
 <ASSIGN>
 <PLAN>\$\$\$AgreeFix
 <STATUS>%%TBC

Referee Comment	In the second paragraph of page 3, it says 'using matched replication timing data in multiple cancer types significantly outperforms an approach in a which one restricts the analysis to replication timing data from the unmatched HeLa-S3 cell line.' This statement is confusing and does Figure 2A or 2B supported it?
Author Response	
Excerpt From Revised Manuscript	

Formatted Table

<ID>REF3.3 – Presentation of the data figure

<TYPE>\$\$\$Presentation
 <ASSIGN>
 <PLAN>\$\$\$AgreeFix
 <STATUS>%%TBC

Referee Comment	In Figure 1, "top tier" should point to cell types that is mentioned in the content. However, we also see SNV, SV, Mutation, etc.
Author Response	
Excerpt From Revised Manuscript	

Formatted Table

<ID>REF3.4 – Regarding enhancer detection algorithm

<TYPE>\$\$\$Presentation

<ASSIGN>

<PLAN>\$\$\$AgreeFix

<STATUS>%%TBC

Referee Comment	What is a single shape algorithm? The authors point to Supplementary data, but there is no definition there either. Do the authors mean the complete graphs or connected components?
Author Response	
Excerpt From Revised Manuscript	

Formatted Table

<ID>REF3.5 – Regression coefficients of BMR

<TYPE>\$\$\$BMR

<ASSIGN>

<PLAN>\$\$\$AgreeFix

<STATUS>%%TBC

Referee Comment	For Figure 2B, what does 'regression coefficients of remaining features' mean? Does that mean beta_0 or the remaining regression noise? From Figure 2B, the coefficient to regression is rounded to -0.001 and 0.001. How should we understand these values? If the coefficients are for the main features, we would be expecting higher coefficients, wouldn't we? In this case, does it mean the lower the better?
Author Response	
Excerpt From Revised Manuscript	

Formatted Table

<ID>REF3.6 – Validation of extended gene

<TYPE>\$\$\$Annotation

<ASSIGN>

<PLAN>&&AgreeFix

<STATUS>%%TBC

Referee Comment	For Figure 2C, more explanation is needed on how to form an extended gene. For the Figure 2D and its description on the third paragraph of page 4 (as well as Figure 3A), did the authors validate all the genes systematically? Is there any validation rate showing the precision rate of the method? Are there any novel oncogenes detected by the method?
Author Response	

Formatted Table

Excerpt From Revised Manuscript	
---------------------------------	--

<ID>REF3.7 – Logic gates

<TYPE>\$\$\$Network

<ASSIGN>

<PLAN>&&&AgreeFix

<STATUS>%%%TBC

Referee Comment	Are circuit gates necessary for Fig 3B? There are OR, AND and NOT gates used. For Figure 3C(i), what is the meaning of the values between the green and yellow dots (MYC and *)? The figure legends are not explaining the figure very well and many details are omitted.
Author Response	
Excerpt From Revised Manuscript	

Formatted Table

<ID>REF3.8 – Network hierarchy

<TYPE>\$\$\$Hierarchy

<ASSIGN>@@@DL

<PLAN>&&&AgreeFix

<STATUS>%%%50DONE

Referee Comment	For Figure 4, what does the star symbol (*) mean in the legend? Did the authors use a different grey color to show
-----------------	--

Formatted Table

	the connection between TFs? I'm not able to read the grey gradient for the edges.
Author Response	We thank referee for point out this issue. We have updated the figure 4 to show the significance testing of network hierarchy analysis. If a p-value is less than 0.05 it is flagged with one star (*). If a p-value is less than 0.01 it is flagged with two stars (**). If a p-value is less than 0.001 it is flagged with three stars (***)
Excerpt From Revised Manuscript	

<ID>REF3.9 – Network rewiring

<TYPE>\$\$\$Network

<ASSIGN>@@@DL

<PLAN>&&&AgreeFix

<STATUS>%%%75DONE

Deleted: >

Deleted: TBC

Referee Comment	For Figure 5B, what does the vertexes and edges represent? I guess they represent genes and their network connection, respectively? How did you select the genes and why are some of them "thick" while others "thin"?
Author Response	We thank referee for pointing this out. First of all, you are correct that vertexes are representing genes and edges are representing regulatory linkage between TFs and genes. We have used colors and thickness to show regulatory rewiring between cell types. Thick edges are shown to highlight rewiring events while thin edges mean gene linkages are retained between cell types.
Excerpt From Revised Manuscript	

Formatted Table

Referee #4 (Remarks to the Author):

<ID>REF4.1 – Strengths of the Paper

<TYPE>\$\$\$NoveltyPos
<ASSIGN>@@@MG,@@@JZ
<PLAN>&&&AgreeFix
<STATUS>%%75DONE

Deleted: <ID>REF4.0 – Preamble .

... [13]

Referee Comment	I fully acknowledge that the manuscript proposes a very important approach from detecting the mutations that are most relevant for each specific type of cancer, integrating epigenome data, transcription factor binding, chromatin looping to focus on key regions: ultimately, this work demonstrates the importance of functional data beyond the primary sequence of the genome. Other important aspects include the comprehensiveness and breadth of the data, the analysis and ultimately the whole integrated approach, which goes beyond commonly seen genomics analysis. However the manuscript is not trivial to read and digest in the first round: anyway I believe that the message, including the importance of the integration multiple types of data, is very important.
Author Response	We thank the referee for the positive comments.

Formatted Table

<ID>REF4.2 – Changing the presentation of the supplement

<TYPE>\$\$\$Text,\$\$\$Presentation
<ASSIGN>@@@DC,@@@JZ
<PLAN>&&&AgreeFix
<STATUS>%%75DONE

Referee Comment	Yet, efforts to make the manuscript more readable will be quite important. For instance, I could understand several sections of the manuscript after reading carefully the not so short supplementary part. The strategy of sample selection was easier to understand after seeing the first figure of the supplementary information, as well as fig S1-3 regarding the number of normal vs cancer cell lines. I'm not sure what the
-----------------	--

Formatted Table

	space limitation for this manuscript will be, but clarity should be an important component of a Nature paper.
Author Response	<p>We thank the referee for pointing out that it is sometimes hard to see the full documentation of our methods in the main part and one has to look at the extensive supplements. We are well aware of this fact. The very large scale of the supplement is typical for large genomic paper. We, in fact, have been actively discussing with Nature Publishing and other companions about the supplement with regard to the main text. We have attempted to put important contents in the supplement and to structure it very carefully.</p> <p>We admit that maybe this construction is not that intuitive. We are prepared to work very hard to make the structure of the supplement understandable. We have tried to revise it to make these clearer and also to move more into the main text, though we think given the current main text limitations of a typical paper in Nature and the scale of the results in the data in this paper, it is not easy to put everything into the main text. We are preparing to work constructively with the referees and the others to make this clear.</p>
Excerpt From Revised Manuscript	[JZ2MG: is there an excerpt here?]

<ID>REF4.3 – Trimming and editing parts of the manuscript

<TYPE>\$\$\$Text,\$\$\$Presentation
 <ASSIGN>@@@DC,@@@JZ
 <PLAN>&&&AgreeFix
 <STATUS>%%%75DONE

Referee Comment	1) The manuscript is quite complex and efforts are needed to improve clarity. Some of the text can seem to be somehow redundant or not needed (for instance, general comments about the ENCODE project; or the Step-Wise prioritization scheme (page7; other parts at page 7, for instance).
Author Response	We thank the referee for his/her suggestions on our presentations. As requested, we have trimmed and edited these sections in our revised manuscript.

Formatted Table

<ID>REF4.4 – [Loss of diversity in cancer cells](#)

<TYPE>\$\$\$CellLine

<ASSIGN>@@@JZ,@@@DL

<PLAN>&&MORE

<STATUS>%%75Done

[\[JZ2MG: I moved the limitation of cells line to the beginning of 4.5. This can change a negative point to a positive point. Please comment this move\]](#)

Referee Comment	I have seen data in other studies, showing that many of cancer cell transcriptome are quite similar to each other, if compared to initial or primary cells, showing that in particular cancer cells lose diversity.
Author Response	We thank referee for bringing this point and we feel it is a good comment. Actually, the referee is correct many of the cancer transcriptome is similar to each other and we made a new figure in our revised version.
Excerpt 1 From Revised Manuscript	One of the strengths of ENCODE release 3 is massive expansion of functional genomic data into various primary cells and tissue types. In this revision, we have extensively explored the chromatin landscape and expression patterns across all of available ENCODE primary cells and tissues, and compared them with existing immortalized cell lines with deep annotations. We have chosen CTCF ChIP-seq and RNA-seq, which has the most abundant number of cell types in ENCODE, as examples to highlight this point. We looked at differential binding patterns of CTCF at promoter regions across cell types. The t-SNE plot of CTCF network shows that most of normal cell lines form a cluster together with healthy primary cells, and cancer cell lines can be linearly separable from their normal counterparts.

Deleted: Comparison

Deleted: tissues to cell lines

Deleted: ,\$\$\$Validation

Deleted: ,@@@Peng

Deleted: 50DONE

Deleted: -

Deleted: 2)

Moved down [6]:One of the limitations of the analysis are the cells that are central in the ENCODE, that are immortalized, including cancer cells and "normal" immortalized counterparts. Most of these cell lines have been kept in culture for decades and further selected for cell growth very extensively. Many of the cell lines may have/have accumulated further mutation and rearrangements, if compared to what cancer cells are at the moment that they leave the human body. The authors accurately acknowledge, in the discussion, stating that it is difficult to match cancer cells with the right normal counterpart; it may also be even more difficult to define what are they really (

Formatted Table

Deleted:)

Deleted: Author ... [14]


Formatted Table

Deleted: We thank referee for bringing this point and we feel it is a good comment. Actually, the referee is correct many of the cancer transcriptome is similar to each other and we made a new figure in our revised version. ... [15]

Formatted: Left

Deleted:


Formatted: Font:Arial



<Figure update candidate: CTCF regulatory networks based on all available ENCODE ChIP-seq shows clustering of stem-like state cell types (Blue). All cancer cell lines (Red) were clustered closer to stem-like cell types than normal cell types (Green).>


Excerpt 2
From
Revised
Manuscript

We performed RCA/PCA analysis on RNA-Seq, shRNA RNA-Seq, and TF ChIP-seq data and found that cancer cells tend to cluster together and stay away from their normal counterparts.



<ID>REF4.5 – Validate the cell line results using tissue data
 <TYPE>\$\$\$CellLine,\$\$\$Validation
 <ASSIGN>@@@JZ,@@@DL,@@@Peng,@@@DC

- Deleted: ... [16]
- Deleted: the leslie data & also some transcriptome analysis (5 genome-wide platforms, incl. RNA-seq, 1 proteomic platform, Hoadley et al. 2014) ... [17]
- Formatted: Font:Arial
- Formatted: Font:Arial
- Deleted: paper (5 genome-wide platforms, incl. RNA-seq, 1 proteomic platform, Hoadley et al. 2014) ... [18]
- Formatted: Font:Arial
- Deleted: the imputed stuff on roadmap tissue to show similar results (5 genome-wide platforms, incl. RNA-seq, 1 proteomic platform, Hoadley et al. 2014) ... [19]
- Formatted: Font:Arial
- Deleted: line as a validation (5 genome-wide platforms, incl. RNA-seq, 1 proteomic platform, Hoadley et al. 2014) ... [20]
- Formatted: Font:Arial
- Deleted: transcriptome http://science.sciencemag.org/content/357/6352/eaan2507 (5 genome-wide platforms, incl. RNA-seq, 1 proteomic platform, Hoadley et al. 2014) ... [21]
- Formatted: Font:Arial
- Deleted: human tissues ... [22]
- Formatted: Font:Arial, 11 pt, Not Highlight
- Deleted: were found to converge into common subtypes.”
- Formatted: Font:Arial
- Deleted: (5 genome-wide platforms, incl. RNA-seq, 1 proteomic platform, Hoadley et al. 2014) ... [23]



Deleted: ... [24]

<PLAN>

<STATUS>%%%100DONE

[JZ2MG: ongoing]

Deleted: 75DONE

Deleted: [JZ2PE: use the cristina leslie ATAC-Seq data set]


Referee Comment	<p><u>One of the limitations of the analysis are the cells that are central in the ENCODE, that are immortalized, including cancer cells and "normal" immortalized counterparts. Most of these cell lines have been kept in culture for decades and further selected for cell growth very extensively. Many of the cell lines may have/have accumulated further mutation and rearrangements, if compared to what cancer cells are at the moment that they leave the human body. The authors accurately acknowledge, in the discussion, stating that it is difficult to match cancer cells with the right normal counterpart; it may also be even more difficult to define what are they really (I have seen data in other studies, showing that many of cancer cell transcriptome are quite similar to each other, if compared to initial or primary cells, showing that in particular cancer cells lose diversity).</u></p> <p><i>It would be appropriate to (computationally) verify at least a small part of the data in other systems, taking from published studies including normal cells control and primary cancers.</i></p>
Author Response	<p>We take the referee's comment to heart and we agree with the reviewer that it is important to verify the discoveries from cell lines from primary cancers.</p> <p>In the revision, we compared the concordance level of our conclusions made from ENCODE cell line data to observations from patients with primary cancers. And we clarified that although ENCODE data are profiled in cell culture models, the regulatory targets are still representative of the gene regulations in human cancers. We have added a new section in the revised supplementary file for more discussions.</p>
Excerpt From Revised Manuscript	<p>We predicted the regulatory activities of transcription factor (TF) MYC using a ChIP-Seq profile in MCF-7 cells. We found that the MYC regulatory activity is highly correlated with the MYC expression across TCGA breast tumors (Supplementary Figure Xa). For most TFs, their regulatory activities predicted using ENCODE ChIP-Seq profile in cell lines are significantly correlated with their expression levels across breast tumors (Supplementary Figure Xb). Moreover, using the same MCF-7 ChIP-Seq profile, the MYC regulatory activity predicted for lung tumors is also significantly correlated with MYC expression level in TCGA lung cancer (Supplementary Figure Xa). These results indicate that the ChIP-Seq profiles from a particular cell line can capture regulatory targets in human tumors</p>

Moved (insertion) [6]

Formatted Table

Formatted: Font:Italic, Underline

from diverse cancer types. To select ChIP-Seq or eCLIP profiles that are representative of the regulatory targets in human cancers, we only reported the results of TFs or RBPs whose regulatory activities are significantly correlated with their gene expression level in each TCGA cohort (Supplementary Figure Xc).




Supplementary Figure X. The clinical relevance of ENCODE cell line data in human primary tumors.

(a) The correlation between *MYC* expression level and regulatory activity across tumors. The *MYC* regulatory activity in each tumor was predicted using the ChIP-Seq profile in MCF-7 cell line. The Pearson correlation between *MYC* gene expression level and regulatory activity were computed across tumors in each cancer type. The statistical significance of Pearson correlation was tested by the two-sided student t-test. BRCA: breast invasive carcinoma. LUSC: lung squamous carcinoma.

(b) The distribution of correlation p -values in TCGA breast cancer. For each TF, we tested the statistical significance of Pearson correlation between TF expression levels and regulatory activities predicted across tumors through two-sides student t tests as panel a. For TCGA breast cancer cohort, most p -values are very significant with a few non-significant values.

The fraction of regulators with statistically significant correlations in different cancer types for ChIP-Seq and eCLIP networks. In each TCGA cancer type, we computed the correlations between regulator expression levels and regulatory activities across tumors for all regulators (TFs, or RBPs). We selected regulators with statistically significant correlations through two-sided student t test (FDR < 0.05).



<ID>REF4.6 – Relationship of H1 to other stem cells

<TYPE>\$\$\$Stemness\$\$\$Calc



<ASSIGN>@@@DL,@@@PE,@@@DC

<PLAN>&&&AgreeFix,&&&MORE

<STATUS>%%%75DONE

Referee Comment	3) One of the conclusions, deriving from the analysis of H1-hESC is the some cancer are "moving away from stemness". However, while it is true that the cancer cells pattern diverge from the H1 cells, H1 is a human embryonic stem cells: although interesting, <u>H1 may not necessarily be the best cells to compare with tumor phenotype</u> . Authors should discuss/defend of further elaborate on this approach. I believe that a key analysis should be done against <u>other stem cells</u> (like tissutal stem cells, etc.).
Author Response	<p>We thank the referees for bringing this point out and we have done what they suggested. We have chosen H1-hESC because it offers the broadest ChIP-seq coverage and has the most amount of other assays in ENCODE. In our revised manuscript, we have expanded our analysis to other stem cells. We have compared other available stem-related cell types, as suggested by the referee, to H1-hESC to show that H1-hESC is not very different from other stem cells from tissues. We have evaluated regulatory activity of all ENCODE biosamples and across all available stem-like cells in ENCODE and measured the distance between stem-like cells. We show that H1-hESC is not far distinct from other stem-like cells. As shown earlier, one analysis we have added is to look at regulatory networks of CTCF, one of the most widely assayed TF in ENCODE. As expected, all of stem-like cell types formed a cluster, suggesting stem-like cell types have a distinct regulatory profile from normal and cancerous cell types, and stem-like cells including H1 and iPSCs have similar regulatory patterns .</p> <p>Another analysis we added was to look at gene expression profiles of all available ENCODE cell types. In agreement with the previous analysis, gene expression profiles of stem-like cell types were very similar to each other and formed a cluster when projected onto 2D RCA space.</p>

Formatted Table



<ID>REF4.7 – Fixes for Figure 1

<TYPE>\$\$\$Presentation,\$\$\$Later

<ASSIGN>@@@DL

<PLAN>&&&AgreeFix

<STATUS>%%%75DONE

Referee Comment	4) I have difficulties to fully understand Fig.1, in particular the patient cohort (PC) at the bottom of the “depth approach” (just above the green box of cell -specific analysis). The two rows are at the bottom of the columns report mutation and expression, but they belong to the columns of the cell lines (K562, HepG2, etc). I just simply do not understand that part of the figure, in particular the relation between cell lines and the patient cohort (the figure legend does not help, and also supplementary material did not help).
Author Response	We thank referee for the suggestion. In the revision we have extensively revised the figure 1. We understand that numbers at the mutation and expression rows can be misleading, so we have separated cohort-based data matrix out of cell-type data matrix. In addition, more emphasis was put into the overview schematic to highlight the value of ENCODEC as a resource.

Formatted Table

Excerpt From Revised Manuscript	
---------------------------------	--

<ID>REF4.8 – SVs affecting BMRs & Network

<TYPE>\$\$\$BMR,\$\$\$Network,\$\$\$Calc
 <ASSIGN>@@@DL,@@@XK, @@@TG,@@@STL
 <PLAN>&&&AgreeFix,&&&MORE
 <STATUS>%%30DONE

[JZ2DL, XM, TG, STL: woould you please help to fill in the stuff?]


Deleted: [JZ2MG: to disc next week] -

Referee Comment	5) The analysis assumes that genomes of all the cells discussed are essentially the same. However, for many of the cancer genomes, there have been rearrangements, often dramatic like Chromothripsis. How is this affecting the BMR and the linking of non-coding elements to the target genes? How many of the cells analyzed were dramatically rearranged?
Author Response	<p>The referee asked us to comment on the relationship of structural variants, BMR, and network wiring. We think these are very good suggestions and we wished we had taken that more in this mission.</p> <p>In the revision, we have definitely taken this comments to heart and have added in main text figures that look at the degree to which structural variants, or SVs, mature background mutational rate, and they also affected the network rewiring. We think this is an ideal illustration of the ENCODE data since, in addition to mapping a lot about the function of the genome, some of the new incurred data sets actually give rise to structural variants meaning that structural variants are an integral output of the product. Relating them to network wiring and background mutation rate is an ideal illustration of the value of the data and the project. We have constructed a number of new main figures that address this and we quite heartly thank the referee for pointing this out. To summarize our conclusion,</p> <p>First, we did observe an elevated SNV/indel rate around the breakpoints. Second, we explored the SV introduced enhancer gain/loss events and relate them to gene expression changes. Third, we studied the relationship of SNVs to network rewirings</p>

Formatted Table



Excerpt 1
From
Revised
Manuscript

Regarding the relationship of SNV to SV



Prudent 1KG filtered: 1) <math><.5</math> reciprocal overlap 2) no bkpt +/- 100bp; N= \sim 5.9k

Black line: window smoothed (window_size=10, step=1)



Deleted:

... [25]

<ID>REF4.9 – Aspects of heterogeneity related to cell lines

<TYPE>\$\$\$CellLine,\$\$\$Text

<ASSIGN>@@@WM,@@@JZ,@@@MRS

<PLAN>&&&AgreeFix

<STATUS>%%50DONE

Referee Comment	6) Most cancers are not necessarily represented by a single cell type used to obtain genomics data in this study, but contains numerous types of cells with different mutations, as well as normal cells, infiltrating cells, all in a three dimensional structure, often producing metastatic colonizing other organs. However, this study focuses only on comparisons between cells. These limitations should be better discussed, also to put in perspective future studies on single cells.
Author Response	We thank the referee for bringing this up and we <u>completely</u> agree with the referee that genomic and epigenomic heterogeneity in tumor cells, as well as heterogeneity in the tumor microenvironment (e.g., immune cell infiltrates, hormonal factors, normal cell populations, etc.) are significant factors in tumor growth and development. This is a limitation of the current technique, which we now discuss with greater emphasis <u>(more details in the excerpt below)</u> .
Excerpt From Revised Manuscript	<u>One limitation of the current ENCODE data is that most of the current release of data is performed over a number of cells. However, genomic and epigenomic heterogeneity in tumor cells, as well as heterogeneity in the tumor microenvironment (e.g., immune cell infiltrates, hormonal factors, normal cell populations, etc.) are significant factors in tumor growth and development. We believe that in the further, the development of single-cell sequencing technologies may capture important tumor biology present and provide new insights in cancer.</u>

Deleted: [JZ2MG: special attention. To disc next ... [26]

Formatted Table

Deleted: totally

Deleted: .

Deleted: - ... [27]

Deleted: We predicted the regulatory activities of transcription factor (TF) MYC using a ChIP-Seq profile in MCF-7 cells. We found that the MYC regulatory activity is highly correlated with the MYC expression across TCGA breast tumors (Supplementary Figure Xa). For most TFs, their regulatory activities predicted using ENCODE ChIP-Seq profile in cell lines are significantly correlated with their expression levels across breast tumors (Supplementary Figure Xb). Moreover, using the same MCF-7 ChIP-Seq profile, the MYC regulatory activity predicted for lung tumors is also significantly correlated with MYC expression level in TCGA lung cancer (Supplementary Figure Xa). These results indicate that the ChIP-Seq profiles from a particular cell line can capture regulatory targets in human tumors from diverse cancer types. To select ChIP-Seq or eCLIP profiles that are representative of the regulatory targets in human cancers, we only reported the results of TFs or RBPs whose regulatory activities are significantly correlated with their gene expression level in each TCGA cohort (Supplementary Figure Xc). - ... [28]

<ID>REF4.10 – lncRNAs and BMR

<TYPE>\$\$\$BMR,\$\$\$Calc
 <ASSIGN>@@@JZ
 <PLAN>&&&AgreeFix
 <STATUS>%%50DONE

Referee Comment	7) When analyzing the BMR in cancer, did the author estimate the mutation rate in the lncRNAs? Is there any other interesting lesson from the analysis of the non-coding regions and their mutations rate?
Author Response	We thank the referee to point out this. We have added the analysis of lncRNA by comparing BMRs in genes and lncRNAs.

Formatted Table

Excerpt From Revised Manuscript	
---------------------------------	--

<ID>REF4.11 – (Minor) updates to figure numbering in supplementary

<TYPE>\$\$\$Minor,\$\$\$Presentation
 <ASSIGN>@@@JZ
 <PLAN>&&&AgreeFix
 <STATUS>%%75DONE

Referee Comment	In the supplementary material, there is room to improve figures (some numbers are too small).
Author Response	We thank the referee to point out this and we have fixed in our revised manuscript
Excerpt From Revised Manuscript	

Formatted Table

<ID>REF4.12 – (Minor) Figure legends


<TYPE>\$\$\$Minor,\$\$\$Presentation
 <ASSIGN>@@@JZ
 <PLAN>&&&AgreeFix
 <STATUS>%%75DONE

Referee Comment	Figure legends. Figure legends are essential but I struggled to understand the figures based on the legends only.
Author	We thank the referee to point out this and we have fixed in our revised manuscript

Formatted Table

|
|
|

Response	
Excerpt From Revised Manuscript	



Referee #5 (Remarks to the Author):

<ID>REF5.0 – Preamble

<TYPE>\$\$\$Text

<ASSIGN>@@@MG,@@@JZ

<PLAN>&&&AgreeFix

<STATUS>%%%75DONE

We would like to appreciate the referee's feedback. We found that many of the suggestions, such as further power analysis, the false positive rate of rewiring, comparison with other networks, cross-validation using external data, are quite valuable and we significantly expanded them in our revised manuscript as suggested. The referee mentioned that, but the novelty of the paper is lacking. We also thank the referee to point out his/her confusion about whether this is prospective or biology paper. We want to make it clear that this paper is to be considered as a "resource" paper, not a novel biology paper. We feel that cancer is the best application to illustrate certain key aspects of ENCODE data and analysis - particularly deep annotations and network changes. We have listed some more details about the novelty of this paper as below.

Contribution	Subtypes	Data types	ENCODE experiments
Processed raw signal tracks	Histone modification	Signal matrix in TSV format	2015 Histone ChIP-seq
	DNase I hypersensitive site (DHS)	Signal matrix in TSV format	564 DNase-seq
	Replication timing (RT)	Signal matrix in TSV format	135 Repli-seq and Repli-ChIP
	TF hotspots	Signal track in bigWig format	1863 TF ChIP-seq
Processed quantification matrix	Gene expression quantification	FPKM matrix in TSV format	329 RNA-seq
	TF/RBP knockdowns and knockouts	FPKM matrix in TSV format	661 RNAi KD + CRISPR-based KO
Integrative annotation	Enhancer	Annotation in BED format	2015 Histone ChIP-seq 564 DNase-seq STARR-seq
	Enhancer-gene linkage	Annotation in BED format	2015 Histone ChIP-seq 329 RNA-seq

Formatted Table

	Extended gene	Annotation in BED format	1863 TF ChIP-seq 167 eCLIP Enhancer-gene linkage
SV and SNV callsets	Cancer cell lines	Variants in VCF format	WGS BioNano Hi-C Repli-seq
Network	RBP proximal network	Network in TSV format	167 eCLIP
	Universal TF-gene proximal network	Network in TSV format	1863 TF ChIP-seq
	Tissue-specific TF-gene proximal network	Network in TSV format	1863 TF ChIP-seq
	Tissue-specific imputed TF-gene proximal network	Network in TSV format	564 DNase-seq
	TF-enhancer-gene network level 1-3	Network in TSV format	2015 Histone ChIP-seq 564 DNase-seq

Specifically for the BMR estimation part, the reviewer mentioned that there had been many existing references focusing on applications like cancer driver detection. First, we thank the referee for pointing out to a lot of related references. On the reference side, we have listed many of the papers as the referee suggested and compared them with our approach. We have acknowledged the efforts of many of these references, and in the revised version we have further expanded our reference list for some the publications after our initial submission date. We want to emphasize that the richness of the ENCODE data can help many of the methods used in these papers. With a larger pool of covariate selection, the estimation accuracy can be improved.

Deleted: significantly

Reference	Initial	Revised	Main point	Comments
Lawrence et al, 2013	Cited	Cited	Introduce replication timing and gene expression as covariates for BMR correction	Replication timing in one cell type
Weinhold et al, 2014	Cited	Cited	One of the first WGS driver detection over large scale cohorts.	Local and global binomial model
Araya et al, 2015	No	Cited	Sub-gene resolution burden analysis on regulatory elements	Fixed annotation on all cancer types
Polak et al (2015)	Cited	cited	Use epigenetic features to predict cell of origin from mutation patterns	Use SVM for cell of origin prediction, not specifically for BMR
Martincorena et al (2017)	No (out after our submission)	Cited	Use 169 epigenetic features to predict gene level BMR	No replication timing data is used
Imielinski (2017)	No	Yes	Use ENCODE A549 Histone and DHS signal for BMR correction	Limited data type used from ENCODE
Tomokova et al. (2017)	No	Yes	8 features (5 from ENCODE) for BMR prediction and mutation/indel hotspot discovery	Expand covariate options from ENCODE data
huster-Böckler and Lehner (2012)	Yes	Yes	Relationship of genomic features with somatic and germline mutation profiles	NOT specifically for BMR
Frigola et al. (2017)	No	Yes	Reduced mutation rate in exons due to differential mismatch repair	NOT specifically for BMR
Sabarathan et al. (2016)	No	Yes	Nucleotide excision repair is impaired by binding of transcription factors to DNA	NOT specifically for BMR
Morganella et al. (2016)	No	Yes	Different mutation exhibit distinct relationships with genomic features	NOT specifically for BMR
Supek and Lehner (2015)	No	Yes	Differential DNA mismatch repair underlies mutation rate variation across the human genome.	NOT specifically for BMR

<ID>REF5.1 – Positive comment of the paper

<TYPE>\$\$\$Text

<ASSIGN>@@@MG,@@@JZ

<PLAN>&&&AgreeFix

<STATUS>%%%DONE

Referee Comment	While the resources provided in this manuscript are potentially interesting for the cancer genomics community and comprise an extensive body of work
-----------------	--

Formatted Table

Author Response	We thank the referee for the positive comment.
-----------------	--

<ID>REF5.2 – BMR: [novelty compared to previous work](#)

<TYPE>\$\$\$Text
 <ASSIGN>@@@JZ
 <PLAN>&&&AgreeFix
 <STATUS>%%%75DONE

Deleted: DONE

Referee Comment	1. The manuscript does not clearly state innovation and novelty over previously published data and methods. Several published studies have used epigenomic data types, including replication time and histone modifications from ENCODE and other sources, to model background mutational background density and define genomic elements of interest. The use of the Negative Binomial/gamma-Poisson distributions to model mutational background in cancer has also been published (Imielinski et al 2016; Martincorena et al, 2017).
-----------------	--

Formatted Table

Author Response	<p>We thank the reviewer for bringing out these references. We want to point out that the Martincorena et al. paper came out in Nov 2017, almost three month after our submission. And it is more focused on positive selection patterns instead of BMR estimation, which makes it unfair for a direct comparison.</p> <p>We also want to clarify that our manuscript is not to claim a new discovery that using matched features are better, but rather to show that the breadth of ENCODE data allows for improved estimates of background mutation rate. We have further acknowledged prior efforts on this topic in our revised manuscript.</p> <p>It is worth to mention that we have released way more genomic features in a ready-to-use format and have shown that it would noticeably improve BMR estimate accuracy if appropriately used. We want to further emphasize two points here.</p> <p>1. ENCODE3 uniformly processed 2017 histone modification data, which makes a much larger pool of features to choose from to potentially improve BMR estimation. Also, the majority of them are actually from real tissues and primary cells (1339 out of 2017).</p>
-----------------	--

Deleted: did notice

Deleted: epigenetic features have been used to estimate

Deleted: and improve driver mutation detection.

Deleted: do

Deleted: intend

Deleted: it is

	2. ENCODE3 provides way more replication timing data. Previously, researchers either use no or only HeLa replication timing for all cancer types (Martincorena et al., 2017, Lawrence et al., 2013), or any of the 16 repli-Seq data from previous ENCODE release. We largely extended this number to 51 cell types (12 cell lines).														
Excerpt From Revised Manuscript	<p>Table S1. Summary of ENCODE3 histone ChIP-Seq data</p> <table border="1"> <thead> <tr> <th>Cell Type</th> <th># histone marks</th> </tr> </thead> <tbody> <tr> <td>tissue</td> <td>818</td> </tr> <tr> <td>primary-cell</td> <td>521</td> </tr> <tr> <td>cell-line</td> <td>339</td> </tr> <tr> <td>in-vitro-differentiated-cells</td> <td>179</td> </tr> <tr> <td>stem-cell</td> <td>114</td> </tr> <tr> <td>induced-pluripotent-stem-cell-line</td> <td>46</td> </tr> </tbody> </table>	Cell Type	# histone marks	tissue	818	primary-cell	521	cell-line	339	in-vitro-differentiated-cells	179	stem-cell	114	induced-pluripotent-stem-cell-line	46
Cell Type	# histone marks														
tissue	818														
primary-cell	521														
cell-line	339														
in-vitro-differentiated-cells	179														
stem-cell	114														
induced-pluripotent-stem-cell-line	46														

Formatted Table

<ID>REF5.3 – [BMR](#): TCGA benchmark

<TYPE>\$\$\$BMR,\$\$\$Calc

<ASSIGN>@@@JZ,@@@WM

<PLAN>&&MORE


<STATUS>%%50DONE,%%CalcDONE


Deleted: 75DONE

Deleted: [JZ2WM: can you please help to paste your stuff here?]

Formatted Table

Referee Comment	2. Throughout, the main manuscript lacks data and statistics supporting the claims made. For example, the performance of tissue-specific background mutation models applied to TCGA data needs to be evaluated against known results and benchmarks from TCGA. It seems that some of these are presented in the extensive supplement and should be moved to the main manuscript.
Author Response	We thank the referee for bringing out this point. We agree that it is important to benchmark the mutation rate estimation. However, we are part of the PCAWG noncoding driver detection group for the joint analysis of TCGA and ICGC data. From our experience in this group, we did not find a gold standard for the whole genome mutation rate estimation. Alternatively, we evaluated the BMR estimation to the commonly used permutation set, which random select a new position within a 50kb window of each somatic variant while preserving the local context.

	<p>1. We applied our mutation driver detection method on the CDS regions of ~20k protein coding regions on the permuted dataset for breast cancer, and found no driver there. QQ plot was added into the supplementary site.</p> <p>2. We down sampled the simulated dataset and xxxx (WM to fill in)</p>
Excerpt 1 From Revised Manuscript	<p>1. QQ plot of the observed vs. uniform p value from Breast cancer permuted data set. Red line is the diagonal line.</p> 
Excerpt 1 From Revised Manuscript	WM downsampling?



<ID>REF5.4 – Improvements of the BMR

<TYPE>\$\$\$BMR,\$\$\$Calc

<ASSIGN>@@@JZ@@@WM

<PLAN>\$\$\$MORE,\$\$\$DisagreeFix,\$\$\$OOS

<STATUS>%%TBC

[\[JZ2MG: only for discuss purpose. I merged this to 5.8. Driver discover is out of scope\]](#)

Referee Comment	<p>3. An improvement of background mutation rate is suggested in the manuscript. But concrete comparisons of discovered drivers with previous work, highlighting how the presented</p>
-----------------	--

Deleted: [JZ2MG: need more advice here? Does it look good?] -

Formatted Table

	approach is more sensitive or improves specificity, are missing.
Author Response	[merged with 5.8, Driver discover is out of scope] Preserve here temporarily for Monday discussion!!!!!!
Excerpt From Revised Manuscript	

Deleted: Part of the previous - ... [29]

<ID>REF5.6 – Power analysis

<TYPE>\$\$\$BMR,\$\$\$Calc

<ASSIGN>@@@JZ

<PLAN>&&&MORE

<STATUS>%%%75DONE

Referee Comment	4. How do the new "compact annotations" lead to improved results over traditional annotations? <u>The power considerations for selecting genomic elements are valuable. "Increased" power of the combined strategy is suggested in the manuscript, yet comparison to prior work is missing.</u>
Author Response	<u>We thank the referee for his/her positive comment on the value of selecting genomic element and suggestion on the power analysis. In our revised manuscript, we expanded our power calculation extensively (see details below) and clearly pointed out the difference of assumptions.</u>
Excerpt 1 From Revised Supplementary file	<u>Regarding compact annotation:</u> <u>In our initial submission, the assumption is that we were trimming off the nonfunctional sites while preserving the functional ones. Two examples can explain the motivation of this assumption.</u> <u>1) Enhancers: Traditionally, enhancers were called as a 1kb peak regions, which admittedly introduced a lot of obviously nonfunctional sites. We believe we can get functional region more accurately by trimming the enhancers down using the exact shapes of many histone marks and further integration with STARR-seq and Hi-C data.</u>

Deleted: [JZ2MG: seems that this referee need to see results not just math equations] -

Moved (insertion) [7]


Deleted: Referee - ... [30]

Formatted Table

Moved (insertion) [8]

Formatted: Font: Times New Roman

2) TFBS hotspots around the promoter region of WDR74. Instead of testing the conventional up to 2.5K promoter region, we can trim the test set to a core set of the promoter region where many TFs bind, which perfectly correlates with the mutation hotspots (red block) for this well-known driver site (blue line for pan-cancer and green line for liver cancer).



[Excerpt 2 From Revised Supplementary file](#)

Following the reviewer's suggestions, in our revised manuscript we show in a formal power analysis that the most important contribution to power comes from including additional functional sites, which is of course by the extended gene concept and then secondarily, from removing non-functional sites, but to a lesser extent. The assumption in our compacting annotations is that we can accurately distinguish the more important functional nucleotides from the less important ones through the guidance of many functional characterization assays.

Admittedly, we are making assumptions and the referee is completely correct in pointing this out. We have tried to be more precise in the text that we are assuming that the large number of ENCODE assays, when integrated, allow us to more directly get the functional nucleotides, but this, of course, is an assumption. It is hard to tell to what degree one can succeed in finding the current events in cancer. It is hard to back this up with the gold standard, but we think that some of the points are self evidently obvious. We have tried to make this clear in text and thank the referee for pointing this out.

Moved up [7]: The power considerations for selecting genomic elements are valuable.

Deleted: 4 .

Moved down [9]: reduced element size need to be added.

Deleted: Again, sensitivity/specificity analyses of driver discovery with large sets, or long vs.

Deleted: Prior efforts to address this problem with restricted hypothesis testing for cancer genes should be cited (Lawrence et al, 2014; Martincorena, 2017) .

Moved up [8]: - ... [32]

Deleted: Author - ... [31]

Deleted: We thank the referee for his/her positive comment on the value of selecting genomic element and suggestion on the power analysis. In our revised manuscript, we expanded our power calculation extensively (see details below). In terms of reference, we cited the Lawrence et al, 2014 paper (and the paper before this one in the same group) in our initial submission and added the Martincorena, 2017, which is published after our submission in Aug 2017. .

Formatted: Font:Times New Roman

Deleted: -

Formatted: Font:Times New Roman

<ID>REF5.7 – [Power analysis; adding more reference](#)


<TYPE>\$\$\$BMR,\$\$\$Text

<ASSIGN>@@@JZ

<PLAN>&&&MORE

<STATUS>%%75DONE

Deleted: Excerpt From -



Deleted: Comparing power

Deleted: to other work

Referee Comment	4. The power considerations ... Prior efforts to address this problem with restricted hypothesis testing for cancer genes should be cited (Lawrence et al, 2014; Martincorena, 2017).
Author Response	We thank the referee for bring out previous efforts. In fact, we cited the Lawrence et al, 2014 paper (and the paper before this one in the same group) in our initial submission. The Martincorena, 2017 was published after our submission for it is impossible for us to cite in the last round. We have added it in our revised manuscript.

Moved (insertion) [10]

<ID>REF5.8 – BMR & Power analysis: detailed driver detection comparison

<TYPE>\$\$\$Power,\$\$\$Text
 <ASSIGN>@@@JZ
 <PLAN>&&&MORE
 <STATUS>%%%25Done

Deleted: TBC

Deleted: [JZ2MG: can we say this is out of scope here? Please advise]

Referee Comment	Again, sensitivity/specificity analyses of driver discovery with large sets, or long vs. <u>reduced element size need to be added.</u> An improvement of background mutation rate is suggested in the manuscript. But concrete comparisons of discovered drivers with previous work, highlighting how the presented approach is more sensitive or improves specificity, are missing.
Author Response	We thank the referee for pointing this out. We want to emphasize that the main goal of our paper is not to make novel driver discoveries but to illustrate that the richness of the ENCODE data can noticeably help the accuracy of BMR estimation. It is out of the scope of our paper to make detailed comparison of cancer driver discoveries. However, we did labeled the known driver genes in our calculations with supporting pubmed IDs. We further compared our results with the PCAWG reports (unpublished data).
Excerpt From Revised Manuscript	To be added by JZ

Formatted Table

Deleted: 5. "Increased" power

Moved (insertion) [9]

Deleted: the combined strategy

Deleted: , yet comparison to prior


Deleted: Following the reviewer's suggestions, we show in a formal power analysis now in the supplement that the most important contribution to power comes from including additional functional sites, this is of course by the extended gene concept and then secondarily, from removing non-functional sites, but to a lesser extent. The assumption in our compacting annotations is that we can accurately distinguish the more important functional nucleotides from the less important ones through the guidance of many functional characterization assays. . . . [34]

<ID>REF5.9 – Annotation: false positive rates of enhancers

<TYPE>\$\$\$Power,\$\$\$Text
 <ASSIGN>@@@JZ,@@@MTG
 <PLAN>&&&AgreeFix
 <STATUS>%%75DONE

Deleted: 8 –

Deleted: DONE


Referee Comment	6. The authors claim that reduction of functional elements increases power to discover recurrently mutated elements. This point needs quantitative support in the main manuscript (some analysis is given in the supplemental). For example, in the enhancer list derived from the ensemble method, what fraction of enhancers are estimated to be false positives?
Author Response	We thank the referee for pointing out the importance of power calculations. As suggested we have added more in both main manuscript and supplementary file (as in the excerpt below) .
Excerpt From Revised Manuscript	<p><u>As for the enhancer part, with the ensemble method, for example, we can get more accurate annotation and pin-point to sequences where transcription factors would actually bind to. To estimate the false positive rate would not be very practical at this stage as there is no gold-standard experiment that could assert an predicted enhancer is definitely negative. Here we took the FANTOM enhancer data set and assess the overlap percentage of our enhancer annotation in each ensemble step. We show that each ensemble step indeed increases the percentage of overlap between our annotation and the FANTOM enhancer set. The overlap percentage for our annotation is much higher than that of the Roadmap annotation, and is also higher than the main encyclopedia enhancer annotation (ccRE).</u></p> 

Formatted Table

Deleted: . . . [35]

Moved (insertion) [11]

Formatted: Font:Times New Roman, Not Italic, No underline



Deleted:

<ID>REF5,10 – Assessing quality of enhancer gene linkage annotation

<TYPE>\$\$\$Annotation,\$\$\$Text

<ASSIGN>@@@KevinYip,@@@SKL

<PLAN>&&MORE

<STATUS>%%%50DONE



Referee Comment	7. The authors claim superior quality of gene-enhancer links and gene communities derived from their machine learning approach. The method should at least be outlined in the main text, and accompanied by data supporting its accuracy and better performance compared to existing approaches.
Author Response	<p>We thank the referee for the comments. In the revised supplementary file, we have added two sections to discuss these points.</p> <p><i>1. Regarding the gene-enhancer linkages</i> ###27mar: to be included from Cao Qin</p> <p><i>2. Regarding the gene community methods</i></p> <p>We have compared the gene community model with other methods like NMF by extending our analysis from 122 GM12878 and K526 dataset to all the 862 TF ChIP-Seq assays included in ENCODE data portal. Analysis showed that our method can better preserve the data structure after dimension reduction.</p>
Excerpt From Revised Manuscript	<p>Mix membership model is a hierarchical Bayesian topic model framework and can help to uncover the underlying semantic structure of a document collection. The core of topic models is Latent Dirichlet Allocation(LDA), which cast the mixed-membership (topics) problem into a hidden variable model of documents. The LDA model has been widely used to analyze a wide variety of data types, including but not limited to text and document data, genotype data, survey and voting data. The advantage of LDA over other algorithms (like SVD, PLSI) used in semantic analysis has been described in Blei 2003.</p> <p>With regards to the referee's question, there is no ready-made answers since the data type (TF target network) and problem-definition of our study are both specific. If we treat the LDA mixed-membership analysis as a dimensionality reduction problem, it is possible to compare how well of a model can reproduce the information of original data, as described in paper (Guo, Y., & Gifford, D. K. (2017). Modular combinatorial binding among human trans-acting factors reveals direct and indirect factor binding. BMC Genomics, 18(1), 45.). The correlations of the original target gene vectors between two TFs are compared with those of dimension reduced vectors. The better method should be much close to original vectors correlations.</p>

Deleted: 9

Formatted Table

Deleted: .

To explore how well the LDA mixed-membership analysis on TF regulatory network, we extend our dataset from 122 GM and K526 samples to all the 862 TF ChIP-Seq assays included in ENCODE data portal. In order to get a reliable correlation, we also increase the number of topic to 50 as the number of TF sample increases. The non-negative matrix factorization (NMF) and Kmeans clustering are used for comparison because the nature of regulatory network requires a non-negative decomposition. The same target dimension $K=50$ was used to NMF and target number of clusters $K=50$ for Kmeans. The Euclidean distance between each data the centroids are used to calculated the correlation. As shown in the figure, the x-axis is original correlation of two TF regulatory target, y-axis is reproduced correlation from LDA document to topic distribution and NMF decomposed matrix. The solid line is the 'loess' smoothing curve for the scattered dots. We can see the LDA method can reproduce the original correlation better than either NMF or Kmeans. Overall correlation between the reproduced pairwise correlation and the original correlation were 0.123 in Kmeans, 0.404 in NMF and 0.788 in LDA.



Deleted:

<ID>REF5.10 – What data sets are used

<TYPE>\$\$\$BMR
<ASSIGN>@@@JZ
<PLAN>&&&Defer
<STATUS>%%%75DONE

Deleted: DONE

Referee Comment	8. From the main manuscript, it is not clear which cancer data sets were analyzed with the new background mutation rate estimates and functional regions. Datasets and sample size should be mentioned explicitly.
Author Response	We thank the referee for bringing out this point. We provide it here in the table and summarized it in a line in the main text.
Excerpt From Revised Manuscript	

Formatted Table

<ID>REF5.11 – Mutational signatures

<TYPE>\$\$\$BMR,\$\$\$Text
<ASSIGN>@@@JZ
<PLAN>&&&AgreeFix
<STATUS>%%%75DONE

Deleted: DONE

Referee Comment	9. Do the authors take into account mutational signatures?
Author Response	We thank the reviewers for pointing this out. In the BMR calculation section, we did consider the local 3mer context effect. But we did not specifically looked into the mutational signatures otherwise. We have made this clear in the discussion section in the revised manuscript.

Formatted Table

Excerpt From Revised Manuscript	We hope that in the future new models that can incorporate, sequence coverage, mutational signatures, small scale features (TF and nucleosome binding), would further integrate the full potential of ENCODE data to better calibrate background mutation rates.
---------------------------------	--

<ID>REF5.13 – Additional QQ plots

<TYPE>\$\$\$BMR,\$\$\$Text
 <ASSIGN>@@@JZ
 <PLAN>&&&AgreeFix
 <STATUS>%%%100DONE

Deleted: 12

Moved (insertion) [12]

Referee Comment	10. The significance analysis of cancer cohorts (Figure 2) should highlight known cancer genes versus those newly found in this study. A QQ-plot should be included to confirm that the algorithm accurately models the background expectation.
Author Response	We thank the reviewers for pointing this out. Yes, we have provided the QQ plot in the supplementary file in our initial submission.

Moved down [13]: <TYPE>\$\$\$BMR,\$\$\$Text - ... [36]

Deleted: DONE - ... [37]

Formatted Table

<ID>REF5.14 – Sequence coverage

<TYPE>\$\$\$BMR,\$\$\$Text
 <ASSIGN>@@@JZ
 <PLAN>&&&AgreeFix
 <STATUS>%%%50DONE

Deleted: 13

Moved up [12]: <TYPE>\$\$\$BMR,\$\$\$Text - ... [38]

Moved (insertion) [13]

Referee Comment	Do the authors include sequence coverage in their method?
Author Response	Thanks for pointing this out. We did not consider coverage but this is a good point. We included in the discussion in our revised manuscript.

Formatted Table


Excerpt From Revised Manuscript	We hope that in the future new models that can incorporate, sequence coverage, mutational signatures, small scale features (TF and nucleosome binding), would further integrate the full potential of ENCODE data to better calibrate background mutation rates.
---------------------------------	--

Moved up [10]: - [39]
 Deleted: 14 – Power analysis for compact annota... [40]


<ID>REF5.15 – BCL6 Questions

<TYPE>\$\$\$Annotation,\$\$\$Calc
 <ASSIGN>@@@XK,@@@TG
 <PLAN>&&&AgreeFix
 <STATUS>%%%.TBC
[\[JZ2JZ: more investigations\]](#)

Deleted: 20Done
 Deleted: [JZ2MG: checking the SV status now, to report next week] -

Referee Comment	11. The authors mention that BCL6 would have been missed in an exclusively coding analysis. In which part of the extended annotations were recurrent BCL6 mutations found? If near the promoter, is the BCL6 5' region a known AID off-target? Are BCL6 mutations in CLL associated with translocations?
Author Response	We thank the referee for this comment. As suggested, we found that there is a mutation hotspot near the first intron of BCL6.
Excerpt From Revised Manuscript	

Formatted Table



<ID>REF5.16 – CHIP-seq vs other computational based networks

<TYPE>\$\$\$Network,\$\$\$Calc
 <ASSIGN>@@@Peng,@@@JZ,@@@DL
 <PLAN> &&&AgreeFix
 <STATUS>%%%.75DONE

Referee Comment	12. The manuscript notes that the new networks presented contain "more accurate and experimentally based" gene links. This claim should be supported with comparisons with existing networks and statistical evaluation. How many of the derived networks are false positives? How many networks are derived in total?
Author Response	<p>We thank the referee for bringing this up this point and we also feel that it is important to make comparison with other existing networks with statistical evaluation. We made the following revisions in the updated manuscript.</p> <p>1. Regarding the proximal regulatory element network:</p> <p><i>1.1 Comparison with Biogrid and String experimental interactions.</i> We showed that the ENCODE ChIP-seq/eCLIP based networks can capture a higher fraction of standard interactions (from manually curated networks from TTRUST) than protein physical networks, including Biogrid and String experimental interactions (see details in excerpt 1).</p> <p><i>1.2 Comparison with DHS-based imputed networks</i> We showed that ENCODE ChIP-seq based networks provided better correlations with DHS-based imputed network provided in Neph et. al. 2012m (see details in excerpt 2).</p> <p><i>1.3 False positive rate estimation of the ChIP-Seq based networks</i> The ENCODE consortium has always enforced a strict data quality standards for all ENCODE produced transcription factor ChIP-seq experiments, which allow us to rigorously control the false positives. (see details in excerpt 3).</p> <p>2. Regarding the distal regulatory element network: With the ChIP-seq, DHS, STARR-seq, ChIA-PET, and Hi-C experiment, ENCODE has a distal TF-enhancer-gene network of high quality, which is less discussed and investigated previously. We feel this is one of the unique aspect of our resource.</p> <p><i>2.1 High quality of enhancer definitions after integrating many histone ChIP-seq and DHS, and STARR-Seq data</i> We provide better enhancer definitions after integrating various assays. Please see details in response to "<ID>REF5.9 – Annotation: false positive rates of enhancers".</p> <p><i>2.2 High quality of enhancer-gene linkages</i> [JZ2JZ: to be added]</p>

Formatted Table

Deleted: below

Deleted: .

Deleted: Seq


Deleted: Seq

Deleted: Here we took the FANTOM enhancer data set and assess the overlap percentage of our enhancer annotation in each ensemble step. We show that each ensemble step indeed increases the percentage of overlap between our annotation and the FANTOM enhancer set. The overlap percentage for our annotation is much higher than that of the Roadmap annotation, and is also higher than the main encyclopedia enhancer annotation (ccRE).


Excerpt 1
From
Revised
Manuscript

Regarding Comparison with Biogrid and String experimental interactions.

To evaluate the quality of ENCODE transcriptional regulatory networks, we utilized the TRRUST database, which manually curated transcriptional regulations from Pubmed articles (Han et al., 2018). We defined the TRRUST interactions as the standard and tested the fraction of standard interactions that other networks can recapitulate. The ENCODE network can capture a higher fraction of standard interactions than protein physical networks, including Biogrid and String experimental interactions (Supplementary Figure X). Moreover, the fraction of standard networks that ENCODE network recapitulated is consistently higher than random. These results supported the higher relevance of ENCODE networks on transcriptional regulation compared to other networks. We also constructed another post-transcriptional network between RBPs and target genes through linking the RBP binding sites on gene 3'UTR regions. To the best of our knowledge, the current study is the first one to study RBP-gene interactions systematically; thus we are not aware of any previous resources that can provide gold standard regulations for comparison.



Supplementary Figure X. ENCODE networks captured a higher fraction of curated regulations than other networks. The TRRUST database manually curated 8,412 transcriptional regulatory interactions from Pubmed articles (Han et al., 2018). We computed the fractions of TTRUST interactions that other networks can recapitulate. Since each ENCODE ChIP-Seq interaction has a regulatory potential (RP) score, we showed the fractions with different RP thresholds. The random fraction for ENCODE network was estimated through 100 perturbed TTRUST networks using the stub-rewiring method that preserved the gene network degrees (Milo et al., 2002).




Deleted:



Excerpt 2
From
Revised
Manuscript

Regarding comparison with imputed network

Our new regulatory network edges are derived from ENCODE TF ChIP-seq experiments, and they provide more accurate gene linkages than imputed networks from other genomic features. To demonstrate the superiority of our new network, we have evaluated our experimentally derived ChIP-seq networks with DHS-based imputed networks from previous publications. We have used two types of ChIP-seq networks. The first one is based

on proximity to TSS and the second one based on target identification from profiles (TIP) method. For imputed network, we used Neph et. al. 2012 (Neph, Shane, et al. "Circuitry and dynamics of human transcription factor regulatory networks." Cell 150.6 (2012): 1274-1286.) TF-to-TF network imputed from DNase I hypersensitive footprints. In addition to Neph et. al. DHS network, we also built our own version of similar DHS network by utilizing the ENCODE DNase-seq dataset. To test the gene linkages, we have utilized ENCODE RNAi based TF knockdown and CRISPR-based TF knockout datasets to test how the target gene linkages defined by various network definition are affected by after KD/KO. Overall, target genes of ENCODE ChIP-seq networks had larger differential expression after knocking down (Supplementary figure X). Moreover, DHS-imputed network derived from ENCODE DNase-seq performed better than the previously published method (not shown here, available in Supplementary document). Supplementary figure X. Evaluation of ENCODEEC network with previously published regulatory network using ENCODE CRISPRi knockdown data. Target genes of ENCODEEC ChIP-seq based networks have larger expression differential after knocking down. Examples of RFX5, SP2, and USF2 shown. More details with full figures comparing all variants of ENCODEEC networks can be found in supplementary document.



	<p style="text-align: center;">K562_CRISPRi_SP2_ENCSR715EDZ</p>  <p style="text-align: center;">K562_CRISPRi_USF2_ENCSR052BWT</p>  <p style="text-align: center;">1</p>
<p>Excerpt 3 From Revised Manuscript</p>	<p><i>Regarding False positive rate estimation of the ChIP-Seq based networks</i></p> <p>In order to ensure that experiments are reproducible, at least two replicates must be performed in either isogenic or anisogenic conditions (For more information about ENCODE 3 ChIP-seq experimental guidelines, please refer https://www.encodeproject.org/documents/ceb172ef-7474-4cd6-bfd2-5e8e6e38592e/@/@/download/attachment/ChIP-seq_ENCODE3_v3.0.pdf).</p>

- Deleted: 2
- Formatted Table
- Formatted: Font:Times New Roman
- Formatted: Font:Times New Roman
- Formatted: Font:Times New Roman
- Formatted: Font:Times New Roman

<p>For transcription factor experiments, 1486 of 1863 (80%) ChIP-seq experiments we have used to compile ENCODEC resources have more than 2 replicates, which allows further quality control of the derived network. ENCODE used IDR (Irreproducible Discovery Rate) framework to ensure reproducibility of high-throughput experiments by measuring consistency between two biological replicates within an experiment. All processed experiments had both rescue and self consistency ratios are less than 2.</p> <table border="1"> <thead> <tr> <th>Self-consistency Ratio</th> <th>Rescue Ratio</th> <th>Resulting Data Status</th> <th>Flag colors</th> </tr> </thead> <tbody> <tr> <td>Less than 2</td> <td>Less than 2</td> <td>Ideal</td> <td>None</td> </tr> <tr> <td>Less than 2</td> <td>Greater than 2</td> <td>Acceptable</td> <td>Yellow</td> </tr> <tr> <td>Greater than 2</td> <td>Less than 2</td> <td>Acceptable</td> <td>Yellow</td> </tr> <tr> <td>Greater than 2</td> <td>Greater than 2</td> <td>Concerning</td> <td>Orange</td> </tr> </tbody> </table> <p>After extensive quality controls for the concordance between replicates, peaks are called using macs2 {"Zhang et al. Model-based Analysis of ChIP-Seq (MACS). <i>Genome Biol</i> (2008) vol. 9 (9) pp. R137"} with p-value cutoff of 0.01.</p>	Self-consistency Ratio	Rescue Ratio	Resulting Data Status	Flag colors	Less than 2	Less than 2	Ideal	None	Less than 2	Greater than 2	Acceptable	Yellow	Greater than 2	Less than 2	Acceptable	Yellow	Greater than 2	Greater than 2	Concerning	Orange
Self-consistency Ratio	Rescue Ratio	Resulting Data Status	Flag colors																	
Less than 2	Less than 2	Ideal	None																	
Less than 2	Greater than 2	Acceptable	Yellow																	
Greater than 2	Less than 2	Acceptable	Yellow																	
Greater than 2	Greater than 2	Concerning	Orange																	

Self-consistency Ratio	Rescue Ratio
Less than 2	Less than 2
Less than 2	Greater than 2
Greater than 2	Less than 2
Greater than 2	Greater than 2

Deleted:

Formatted: Font:Times New Roman

Formatted: Font:Times New Roman

Formatted: Font:Times New Roman

Formatted: Font:Times New Roman

Deleted: Excerpt 3 From .

Regarding quality of enhancers .

Excerpt 4 From .

... [41]

Moved up [11]: As for the enhancer part, with the ensemble method, for example, we can get more accurate annotation and pin-point to sequences where transcription factors would actually bind to. To estimate the false positive rate would not be very practical at this stage as there is no gold-standard experiment that could assert an predicted enhancer is definitely negative. Here we took the FANTOM enhancer data set and assess the overlap percentage of our enhancer annotation in each ensemble step. We show that each ensemble step indeed increases the percentage of overlap between our annotation and the FANTOM enhancer set. The overlap percentage for our annotation is much higher than that of the Roadmap annotation, and is also higher than the main encyclopedia enhancer annotation (ccRE).

Formatted Table

Formatted: Font:Times New Roman

<ID>REF5.17 – MYC KD [Validation](#)

<TYPE>\$\$\$Network,\$\$\$Text



<ASSIGN>@@@DC

<PLAN>&&&AgreeFix


<STATUS>%%%100DONE

Referee Comment	13. MYC is known to have profound effects on gene networks. Have the authors considered comparing the results from their MCF7 knockdown experiment to existing data from similar MYC knockdowns to validate the behavior of the network?
Author Response	We thank the referee for this suggestion and we feel this is a good comment. As suggested we searched for external dataset from multiple platform and cell types and used them to compare with our discoveries. Both datasets confirmed our claims.
Excerpt From Revised Manuscript	1. We carried out these analyses after first identifying an alternative dataset. Specifically, we identified a dataset of gene expression for both MYC knockdowns (as well as a corresponding control) in Gene Expression Omnibus (GEO accession number GSE86504). For these alternative data, gene expression was measured by RNA-seq in the HT1080 cell line. We note that, even though these alternative analyses were conducted on a different cell line, the results we obtain (shown below in the right panels, and now made available in the supplementary materials) validate the behavior of the network, and they are consistent with our previous results (in which gene expression was measured in the MCF-

7 cell line). These comparable results in an alternative cell line suggests that these results are robust.





We also found another array based MYC knockdown data the results correlate well with our discoveries.



Deleted: target nonTarget [42]

Formatted: Font:Times New Roman



<ID>REF5.18 – SUB1 analysis

<TYPE>\$\$\$NoveltyPos,\$\$\$Calc

<ASSIGN>@@@MRS,@@@JL,@@@YY

<PLAN>&&&MORE

<STATUS>%%85DONE

Referee Comment	14. SUB1 is a potentially interesting new cancer gene. The authors should further explore the biology of this gene.
Author Response	<p>We thank the referees for the positive comments. We did follow up with SUB1 in this round of revision.</p> <ol style="list-style-type: none"> We checked SUB1 regulation potential in different cancer types and found that they are consistent as below. We also found that SUB1 tends to bind to the 3UTRs to stabilize its target mRNA. The decay rate of SUB1 is slower than non-targets (p value=1.91e-10). We checked the 3' UTR expression level of SUB1 target genes and found that the target genes are significantly down-regulated upon SUB1 KD. In addition, we found enrichment of SUB1 target genes for CGC (Cancer Gene Census) genes.

Deleted: [JZ2Peng: write something about sub1 decay rate] .


Formatted Table

Formatted: Font:Times New Roman

Formatted: Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"

Formatted: Justified, Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"



Excerpt 1
From
Revised
Manuscript



Inference of RNA binding proteins that drive tumor specific expression patterns. Based on ENCODE eCLIP data, we applied RABIT framework to identify RNA binding proteins (RBP), whose target genes are differentially regulated in diverse TCGA cancer types. (A) For each RBP, the percentage of patients with target genes significantly up regulated (red), down regulated (blue) or not regulated (white) is shown for each cancer type. (B) Hierarchically clustered heatmap was used to show the percentage of patients in each cancer type with RBP target significantly up regulated (red) or down regulated (blue). (C) All TCGA Liver Hepatocellular Carcinoma (LIHC) lung adenocarcinoma (LUAD) patients are divided to two groups according to the *SUB1* activity predicted by RABIT. The overall survival was shown in each group by KM plot. The association between RABIT regulatory activity and overall survival was tested CoxPH regression. (D) The cumulative distributions of gene expression after *SUB1* knock down in HepG2 cell are shown for predicted target genes and non-target genes. The comparison between two categories of expression changes is done through Wilcoxon rank-sum test. (E) The mRNA decay rates are compared between predicted *SUB1* targets and non-target genes as part D.


Excerpt 2
From
Revised
Manuscript

Comparison




Inference of RNA binding proteins that drive tumor specific expression patterns. Based on ENCODE eCLIP data, we applied RABIT framework to identify RNA binding proteins (RBP), whose target genes are differentially regulated in diverse TCGA cancer types. (A) For each RBP, the percentage of patients with target genes significantly up regulated (red), down regulated (blue) or not regulated (white) is shown for each cancer type. (B) Hierarchically clustered heatmap was used to show the percentage of patients in each cancer type with RBP target significantly up regulated (red) or down regulated (blue). (C) All TCGA Lung Adenocarcinoma (LUAD) patients are divided to two groups according to the *SUB1* activity predicted by RABIT. The overall survival was shown in each group by KM plot. The association between RABIT regulatory activity and overall survival was tested CoxPH regression. (D) The cumulative distributions of gene expression after *SUB1* knock down in HepG2 cell are shown for predicted target genes and non-target genes. The comparison between two categories of expression changes is done through Wilcoxon rank-sum test. (E) The mRNA decay rates are compared between predicted *SUB1* targets and non-target genes as part D.

Deleted:



Deleted:



Here we show some IGV examples together with SUB1 binding sites on the 3' UTRs.

Gene	Functions	PMID	Expression profiles of the 3' UTR
BRCA1	The gene is involved in maintaining genomic stability	12677558, 17416853, 23620175, 16551709	
POLE	The gene is involved in DNA repair and replication	26133394, 28423643	
FEN1	The gene is involved in DNA repair and replication	20929870, 22586102	

Formatted: Font:Times New Roman

Gene	Functions	PMID	Expression profiles of the 3' UTR
BRCA1	The gene is involved in maintaining genomic stability	12677558, 17416853, 23620175, 16551709	
POLE	The gene is involved in DNA repair and replication	26133394, 28423643	
FEN1	The gene is involved in DNA repair and replication	20929870, 22586102	

Deleted:

<ID>REF5.19 – Significance of regulatory network hierarchy

<TYPE>\$\$\$Network,\$\$\$Calc

<ASSIGN>@@@DL

<PLAN>&&&AgreeFix




<STATUS>%%100DONE

Referee Comment	15. The manuscript claims that transcription factors placed at the top level of the network hierarchy are enriched in cancer-associated genes and drive expression changes. Both claims need to be supported with statistical tests.
Author Response	We thank the referees for the positive comments. We've done a statistical significance test as requested. The right panel of Figure 4 shows results from Wilcoxon signed-rank test. If a p-value is less than 0.05 it is flagged with one star (*). If a p-value is less than 0.01 it is flagged with two stars (**). If a p-value is less than 0.001 it is flagged with three stars (***). We find that the top-level of the generalized network was enriched with cancer-related TFs with p-value XXX and had larger correlation to drive target gene expression change (p-value XXX).


Formatted Table

Excerpt
From
Revised
Manuscript

Supplementary Figure X.



Deleted:



<ID>REF5.20 – Rewiring of regulatory network

<TYPE>\$\$\$Network,\$\$\$Calc

<ASSIGN>@@@DL

<PLAN>&&&AgreeFix

<STATUS>%%100DONE

Referee Comment	16. In the tumor-normal network comparison, is the fraction of edge changes related to the total number of edges for a given TF? This analysis should further clearly state its null hypothesis (what changes are expected?). What happens when edges are randomly permuted?
Author Response	We thank referee for pointing out this issue. We agree with the referee that we need to be more clear about the rewiring of regulatory network in the revised manuscript.
Excerpt From Revised Manuscript	<p><u>We would like to clarify that the rewiring index is based on the fraction of regulatory edge changes between two cellular contexts. The rewiring index is also normalized across all regulatory proteins, and the sign reflects the direction of rewiring. Details of rScore derivation can be found in Supplementary 5.3. Given this, we assume a null hypothesis to be no change in regulatory edge across cell types. We expect no or minimal change in edges when two cellular contexts are similar. To demonstrate, we selected all available GM12878 ChIP-seq experiments that have at least two replicates, and we performed the same rewiring analysis between isogenic replicates of the same cellular context. The edge changes between two networks will be simply a noise from ChIP-seq experiments.</u></p> <p><u>As expected, when two cellular context are similar, as shown in “baseline”, minimal number of edges do change targets. However, in “rewiring”, TF do change targets extensively when compared across cancerous (K562) to normal (GM12878) cell lines.</u></p>

Formatted Table

Deleted: -

... [44]


Moved down [14]: We would like to clarify that the rewiring index is based on the fraction of regulatory edge changes between two cellular contexts. The rewiring index is also normalized across all regulatory proteins, and the sign reflects the direction of rewiring. Details of rScore derivation can be found in Supplementary 5.3. Given this, we assume a null hypothesis to be no change in regulatory edge across cell types. We expect no or minimal change in edges when two cellular contexts are similar. To demonstrate, we selected all available GM12878 ChIP-seq experiments that have at least two replicates, and we performed the same rewiring analysis between isogenic replicates of the same cellular context. The edge changes between two networks will be simply a noise from ChIP-seq experiments. -

... [45]

Formatted: Font:Times New Roman


Moved (insertion) [14]

Formatted: Font:Times New Roman



Deleted:

... [46]



<ID>REF5.21 – Rewiring analysis in the stem cells

<TYPE>\$\$\$Stemness,\$\$\$Calc
 <ASSIGN>@@@DL,@@@JZ
 <PLAN>&&&AgreeFix
 <STATUS>%%%50DONE

Deleted: TBC

Referee Comment	17. The network change comparisons with the H1 stem cell models need statistical testing for significance. What fraction of the rewired edges are expected to be false positives?
Author Response	<p>We thank referee for the pointing this out. We took the referee's suggestion to heart and significantly expanded the rewiring and H1 stemness part.</p> <p>1. For the false positive question, the ENCODE consortium has always enforced a strict data quality standards for all ENCODE produced transcription factor ChIP-seq experiments, which allow us to rigorously control the false positives. Please refer to Excerpt 3 in response to “<ID>REF5.16 – ChIP-seq vs other computational based networks”.</p> <p>2. We extended our analysis of H1 to RNA-Seq, TF ChIP-Seq (proximal and distal), and TF knockdown data (details in the Excerpt below).</p>

Formatted Table

Deleted: we now have added

Formatted: Justified


Deleted: statistical significance testing

Deleted: H1 stem cell model

Deleted: the revised manuscript.

Excerpt From Revised Manuscript

We performed RCA/PCA analysis on RNA-Seq, shRNA RNA-Seq, and TF ChIP-seq data and found that cancer cells demonstrate a consistent pattern to be more similar to stem cells, as compared to their primary cells of origin.



REF5.22 – Selection of regions for validation testing

Validation, \$\$\$Text
 @@@JZ, @@@DL
 &&AgreeFix
 %%%75DONE

Referee Comment	18. How were the eight regions that were tested functionally selected? Where are these regions located in the genome, and with respect to neighboring genes? How many replicates were performed? What are the p-values?
Author Response	We thank the referee for pointing this out. The eight regions were selected from our integrative promoter and enhancer regulatory elements in MCF-7 cell lines. We prioritized these regulatory regions based on motif breaking power as described in section 6.1 S (see excerpt 1 below). We also provided similar figure for all the other regions in the supplementary file (see excerpt 1 below).
Excerpt 1 From Revised Manuscript	We selected top ten regions with the highest motif breaking power and then tested their regulatory activities using luciferase assay as described in section 6.2 S. Two of ten regions we tested were failed due to issues with plasmid isolation. There were two biological

Formatted Table

Moved (insertion) [15]

Moved up [15]: - ... [48]

Deleted: Author - ... [47]

Formatted Table


Deleted: We thank the referee for pointing this out. We had some of the details in the supplementary but they weren't that well spelled out . We've redone supplementary section 6 and to answer this question. -

Formatted: Font:Times New Roman

Deleted: regions in MCF-7 cell lines. We prioritized these regulatory regions based on motif breaking power as described in section 6.1 S.

Deleted: 3

replicates and three technical replicates for each biological replicate in designing luciferase assays validations. Error bar is representing 95% confidence interval across replicates.




Formatted: Font:Times New Roman


Deleted: mutant and control experiments

Formatted: Font:Times New Roman

Excerpt 2
From
Revised
Manuscript

Details for all tested regions.





<ID>REF5.23 – Presentation and revision to manuscript

<TYPE>\$\$\$Minor,\$\$\$Presentation,\$\$\$Text

<ASSIGN>

<PLAN>&&&AgreeFix

<STATUS>%%TBC

Referee Comment	19. The authors should consider moving the general overview diagrams that constitute much of the main figures to the supplement, and in turn present data-rich figures from there with the main manuscript.
Author Response	<p>We thank for the referee for this comments.</p> <p>We have tried to revise the figures as requested</p> <p>We have fixed figure XX & YY.</p>
Excerpt From Revised Manuscript	

Formatted Table

Formatted: Font:Helvetica Neue

<ID>REF5.24 – Difference between ENCODEC and existing prioritization methods

<TYPE>\$\$\$Validation,\$\$\$Text

<ASSIGN>

<PLAN>&&&AgreeFix

<STATUS>%%100Done

Referee Comment	20. It is not clear how variant prioritization differs or exceeds the variant prioritization method FunSeq published by the same group. Are they complementary approaches?
Author Response	We thank the referee to bring this up. We believe that the method that we used here is new and novel. The important aspect is that it takes advantage of many new ENCODE data and integrates over many different aspects. In particular, it takes into account the STARR-Seq data, the connections from Hi-C, the better background mutation rates, and the network wiring data, which is only possible in the context of the highly integrated and their data available on certain cell lines. We are showing this as an example of the best we can do with this level of integration. The fact that we coupled this with quite successful validation that we believe points to the great value of the integrated incurred data.

Formatted Table

<ID>REF5.25 – [Minor: BMR: provide q-values](#)

<TYPE>\$\$\$Minor,\$\$\$BMR

<ASSIGN>@@@JZ

<PLAN>&&&AgreeFix

<STATUS>%%100DONE

Referee Comment	21. When the authors describe recurrent events, are these significant? If so, please provide p-values (and q-values, when applicable).
Author Response	We thank the referee to point this out. We have the values and q-values all deposited into our online resource and supplementary files. We have made this clearer in our revised manuscript.

Formatted Table

Deleted: BMR

<ID>REF5.26 – [Minor](#): Citation of previous work

<TYPE>\$\$\$Minor,\$\$\$Presentation
<ASSIGN>
<PLAN>&&&AgreeFix
<STATUS>%%100DONE

Referee Comment	22. Prior work using ENCODE chromatin data to define regulatory regions and gene enhancers links should be cited (referred to in the manuscript as "Traditional methods").
Author Response	We thank the referee to point this out. References have been added in the new submission.

Formatted Table

<ID>REF5.27 – [Minor](#): Tumor normal comparison and composite model

<TYPE>\$\$\$Minor,\$\$\$CellLine
<ASSIGN>
<PLAN>&&&AgreeFix
<STATUS>%%100DONE

Referee Comment	23. The use of a "composite normal" is not optimal for tissue or tumor-type specific analyses that the authors advocate. Although the described data resource (ENCODE) may not provide normal control data, normal tissue data from the Roadmap Epigenomics could be included instead (or in addition) to improve the quality of the tumor-normal comparisons.
Author Response	We thank the referee for bringing this out. We did noticed the Roadmap data. Actually, in the new release, ENCODE3 reprocess the complete set of roadmap data and we did include that in our data tables (Figure 1 and supplementary table xxx).
Excerpt From Revised Manuscript	We highlighted the normal tissue data from the Roadmap (processed by ENCODE3) in our revised figure 1 as below.

Formatted Table

<ID>REF5.28 – [Minor](#): Use of H1 for stemness calculation

<TYPE>\$\$\$Minor,\$\$\$Stemness
 <ASSIGN>
 <PLAN>&&&AgreeFix
 <STATUS>%%50DONE

Referee Comment	24. The authors use the H1 embryonic stem cell line as model for “stemness” in cancer. Tumor “stemness” often resembles tissue progenitors, not embryonic stem cells. In the absence of reliable data for such progenitors the authors should note this caveat with their analysis.
Author Response	<p>We thank the referees for bringing this point out. We mainly have chosen H1-hESC because it offers the broadest TF ChIP-seq coverage and also one of the top-tier cell lines with most variety of experimental assays in ENCODE. We agree with the referee that the use of H1 embryonic stem cell line for measuring “stemness” should be further discussed. We, therefore, have revised the manuscript with two additional analysis to show that use of H1-hESC maybe a suitable substitute for a such analysis, especially in the absence of the proper progenitor cell data.</p> <p>1.We first aimed to evaluate regulatory networks of all ENCODE biosamples including many available stem-like cells and profile their differences. We show that H1-hESC is not far distinct from other stem-like cells, and it is a good representation of stem-like state. (see details in Excerpt 1 below).</p> <p>2. We also looked at gene expression profiles of all available ENCODE cell types. In agreement with the previous analysis, gene expression profiles of stem-like cell types were very similar to each other and formed a cluster when projected onto 2D RCA (reference component analysis) space. Tumor cells actually more similar to stem cells, as compared to their normal counterpart (see details in Excerpt 2 below).</p>
Excerpt 1 From Revised Manuscript	We used a regulatory networks of CTCF, one of the most widely assayed TF in ENCODE, to examine their regulatory patterns across different cell types. As expected, all of stem-like cell types formed a cluster, suggesting stem-like cell types have a distinct regulatory profile from normal and cancerous cell types, and stem-like cells including H1 and iPSCs have similar regulatory patterns.


Formatted Table

Deleted: -
 Second analysis we added was to look


Moved down [16]: We used a regulatory networks of CTCF, one of the most widely assayed TF in ENCODE, to examine their regulatory patterns across different cell types. As expected, all of stem-like cell types formed a cluster, suggesting stem-like cell types have a distinct regulatory profile from normal and cancerous cell types, and stem-like cells including H1 and iPSCs have similar regulatory patterns.

Formatted: Font:Times New Roman
 Deleted: Second analysis we added was to look

Deleted: -
 Moved (insertion) [16]
 Formatted: Font:Times New Roman



t-SNE: CTCF




<Figure update candidate: CTCF regulatory networks based on all available ENCODE ChIP-seq shows clustering of stem-like state cell types (Blue). Promoter network of CTCF was projected onto 2D space using t-SNE. All cancer cell lines (Red) were clustered closer to stem-like cell types than normal cell types (Green).>

Deleted: ... [49]

[Excerpt 2 From Revised Manuscript](#)

[Supplementary figure xx: Gene expression profiles of all available ENCODE RNA-seq experiments show that all stem-like cell types form a cluster \(Blue\). Gene expression quantifications were projected onto 2D space using reference component analysis.](#)

PCA of cell clusters in RCA space



<ID>REF5.29 – [Minor](#): Validation of prioritized element

<TYPE>\$\$\$Minor,\$\$\$Validation

<ASSIGN>@@@DL

<PLAN>&&&AgreeFix

<STATUS>%%%75DONE

Referee Comment	25. P-values should be given in Figure 6B for the luciferase reporter assay. The authors may also want to explain why candidate 5, rather than candidate 4 with a much larger expression fold difference was chosen for follow-up.
Author Response	<p>We thank the referee for this comment. We now have added more details of how the validation of candidate regions we selected into the revised supplementary information (please see Excerpt 2 in response to <ID>REF5.22 – Selection of regions for validation testing).</p> <p>The reason we selected the candidate 5 instead of candidate 4 is that the candidate 5 had stronger motif breaking score when disrupted, had higher density of TF binding events, and aligned better with our integrative regulatory region calls.</p> <p>However, we feel that all other regions we tested are among the top prioritized regions and it is important to show these examples. In the revised manuscript, we have also included supplementary plots for all candidate regions tested in details, showing location of neighboring genes, cohort SNV data, histone marks and DHS signal tracks.</p>
Excerpt From Revised Manuscript	Please see figures in Excerpt 2 in response “to <ID>REF5.22 – Selection of regions for validation testing”

Deleted: >

Deleted: [JZ2DL: could you please help to add the tracks? Reason for this?] .

Formatted Table

Deleted: all the

Deleted: tested

Deleted: file.

Deleted: the highest scored variants in our analysis. We made this more clear

Deleted: our new version

<ID>REF5.30 – [Minor](#): SYCP2 and beyond

<TYPE>\$\$\$Minor,\$\$\$NoveltyPos

<ASSIGN>

<PLAN>&&&AgreeFix

<STATUS>%%%TBC

[JZ2JL: can you please do this quickly?]

Referee Comment	26. The discovery of a previously unknown enhancer of SYCP2 is interesting. The authors should consider following up on this lead by integrating existing mutation and expression data from additional studies (e.g. 560 ICGC breast cancers from Nik-Zainal et al).
Author Response	TBC: add this quickly on Monday
Excerpt From Revised Manuscript	

Formatted Table

<ID>REF5.31 – [Minor](#): Utility of ENCODEC

<TYPE>\$\$\$Minor,\$\$\$Presentation

<ASSIGN>

<PLAN>&&&AgreeFix

<STATUS>%%TBC

[JZ2MG: is it OK for the text?]

Referee Comment	27. The abstract mentions the usefulness of ENCODE data for interpretation of non-coding recurrent variants, yet this point is not explored much in the manuscript.
Author Response	<p>We thank the referee for this comment. Actually, we tried to show in Fig 6 how each data type has been integrated to evaluate the function of variants. For example, the histone ChIP-seq, STARR-Seq, and DHS data helped to define function of surrounding element. The histone ChIP-seq, Replication timing, and Expression data help to calibrate local BMR to evaluate mutation rate and somatic burden. TF ChIP-seq/eCLIP data can help to investigate the local nucleotide effect. And Hi-C and ChIA-pet data can help to link noncoding variants to surrounding genes for better interpretation.</p> <p>We made this more clear in our revised manuscript.</p>

Formatted Table

Excerpt From Revised Manuscript	
---------------------------------	--

<ID>REF5.32 – [Minor](#): P-value of survival analysis

<TYPE>\$\$\$Minor,\$\$\$Presentation
 <ASSIGN>@@@DL
 <PLAN>&&&AgreeFix
 <STATUS>%%%75DONE

Referee Comment	28. In Figure 2e, a p-value should be given with the analysis.
Author Response	We thank referee for the comment. We now have updated figure 2e with p-value.
Excerpt From Revised Manuscript	

Formatted Table

<ID>REF5.33 – [Minor](#): Q-value of extended gene analysis

<TYPE>\$\$\$Minor,\$\$\$Presentation
 <ASSIGN>
 <PLAN>&&&AgreeFix
 <STATUS>%%%[75DONE](#)

Referee Comment	29. Figure 2d, q-values should be given for each identified driver gene.
Author Response	We thank referee for the suggestion. We would like to first point out that we were not focused in finding cancer drivers in this analysis. Figure 2d is to illustrate the utility of extended gene. However, we do agree with the referee that adding q-

Deleted: Done

Formatted Table

	value to the figure would be important, so we have updated the figure in the revised manuscript.
Excerpt From Revised Manuscript	

<ID>REF5.34 – [Minor](#): Presentation issue with network hierarchy

<TYPE>\$\$\$Minor,\$\$\$Presentation
 <ASSIGN>
 <PLAN>&&&AgreeFix
 <STATUS>%%100DONE

Referee Comment	30. Figure 4 would benefit from labeling of the network tiers.
Author Response	We thank reviewer for the comment. We fixed the labeling of the network tiers in the revised manuscript.
Excerpt From Revised Manuscript	

Formatted Table

<ID>REF5.35 – [Minor](#): Presentation

<TYPE>\$\$\$Minor,\$\$\$Presentation
 <ASSIGN>@@@DL
 <PLAN>&&&AgreeFix
 <STATUS>%%75DONE

Referee Comment	31. In Figure 6b, it should be clarified whether "samples" refers to genomic locations, patients, or cell lines. The
-----------------	--

Formatted Table

	number of replicates for each experiment should be shown, and p-values between wt and mutant readings should be given.
Author Response	We thank referee for pointing this issue out. We refer "samples" to the genomic locations in the submitted manuscript. We agree with the referee that this could be confusing. We have updated the figure in the revised manuscript.
Excerpt From Revised Manuscript	

<ID>REF5.36 – [Minor](#): Supplementary document

<TYPE>\$\$\$Minor,\$\$\$Presentation

<ASSIGN>

<PLAN>&&&AgreeFix

<STATUS>%%75DONE

Referee Comment	32. The supplement contains multiple reference errors.
Author Response	We thank the referee on this comment and we have made numerous improvements to the supplementary document.
Excerpt From Revised Manuscript	

Formatted Table

Excerpt
From
Revised
Manuscript

in our initial submission.

It is worth mentioning that

help BMR estimation. We also want to emphasize that two points here.

First, To select a perfect “matching” feature (no matter from matter tissue or cell line) is a non-trivial problem due to the heterogeneity of cancer. Even in the Polak 2015 paper, H3K9me3 from Breast luminal epithelial cells is a significant feature in 5 out of cancer types they investigated (Fig. 2a). The noticeably larger pool of functional characterization data from ENCODE3 can actually help to find a matching issue, especially for cancers types that cannot find an obvious “matching” feature from the Roadmap, such as prostate cancer.

Second, the goal of the Polak 2015 paper is to predict the cell of origin, while we are aiming to improve the BMR estimation accuracy. The fact that “matched” cell type features performs better in predicting

Excerpt
From
Revised
Manuscript

[JZ2DS: would you please add xx xx? Also add some text on the CTCF plot]

Cell Type	# histone marks
tissue	818
primary-cell	521
cell-line	339
in-vitro-differentiated-cells	179
stem-cell	114
induced-pluripotent-stem-cell-line	46

Page 15: [7] Deleted

Author

4/1/18 9:22:00 PM

Page 19: [8] Deleted


Author


4/1/18 9:22:00 PM

Page 20: [9] Deleted

Author

4/1/18 9:22:00 PM





Page 22: [10] Deleted Author 4/1/18 9:22:00 PM

Page 25: [11] Moved to page 26 (Move #5) Author 4/1/18 9:22:00 PM

1) Enhancers: Traditionally, enhancers were called as a 1kb peak regions, which admittedly introduced a lot of obviously nonfunctional sites. We believe we can get functional region more accurately by trimming the enhancers down using the exact shapes of many histone marks and further integration with STARR-seq and Hi-C data.

2) TFBS hotspots around the promoter region of WDR74. Instead of testing the conventional up to 2.5K promoter region, we can trim the test set to a core set of the promoter region where many TFs bind, which perfectly correlates with the mutation hotspots (red block) for this well-known driver site (blue line for pan-cancer and green line for liver cancer).

Page 30: [12] Deleted Author 4/1/18 9:22:00 PM

<ID>REF4.0 – Preamble**<TYPE>\$\$\$Text****<ASSIGN>@@@MG,@@@JZ****<PLAN>&&&AgreeFix****<STATUS>%%75DONE**

We would like to appreciate the referee's feedback and positive comments about our resource. We found that many of the suggestions, such as further power analysis, stemness and rewiring, comparison of cell line vs. tissue, cross validation using primary cancer data, are quite valuable. As suggested, we have significantly expanded them while preserving our original goal in our revised manuscript.

Author

Response

We thank referee for bringing this point and we feel it is a good comment. Actually, the referee is correct many of the cancer transcriptome is similar to each other and we made a new figure in our revised version.

23 mar ongoing stuff

####7mar - get pe to do this timputed

the leslie data & also some transcriptome analysis

####7mar either for imputed network OR for the transcription, we take the referee's comment to heart & try to do they we as the the ref suggested

Take one TF from the imputed network

Ask PE on tumor data ATAC

paper

Try to use some

the imputed stuff on roadmap tissue to show similar results

Let peng to use PE's network, compare results?

To use the imputed network in tissue and used the KD data in

Page 41: [20] Deleted **Author** **4/1/18 9:22:00 PM**

line as a validation

KD in tissue external data

**** we've really made better use of the encode knockdown data and highlight
&&&& & knockdowns

PDM references

A pathology atlas of the human

Page 41: [21] Deleted **Author** **4/1/18 9:22:00 PM**

transcriptome

<http://science.sciencemag.org/content/357/6352/eaan2507>

“analyses revealed that gene expression of individual tumors within a particular cancer varied considerably and could exceed the variation observed between distinct cancer types.” (RNA-seq, Uhlen et al. 2017)

Human cancers overexpress genes that are specific to a variety of

Page 41: [22] Deleted **Author** **4/1/18 9:22:00 PM**

human tissues

<http://www.pnas.org/content/102/51/18556>

“The results indicate that many genes that are overexpressed in human cancer cells are specific to a variety of normal tissues, including normal tissues other than those from which the cancer originated.” (microarray, Lotem et al. 2005)

Multiplatform analysis of 12 cancer types reveals molecular classification within and across tissues of origin.

<https://www.ncbi.nlm.nih.gov/pubmed/25109877>

“Five subtypes were nearly identical to their tissue-of-origin counterparts, but several distinct cancer


Page 41: [23] Deleted **Author** **4/1/18 9:22:00 PM**

(5 genome-wide platforms, incl. RNA-seq, 1 proteomic platform, Hoadley et al. 2014)


###

Page 41: [24] Deleted **Author** **4/1/18 9:22:00 PM**


t-SNE: CTCF



<Figure update candidate: CTCF regulatory networks based on all available ENCODE ChIP-seq shows clustering of stem-like state cell types (Blue). All cancer cell lines (Red) were clustered closer to stem-like cell types than normal cell types (Green).>



Prudent 1KG filtered: 1) <.5 reciprocal overlap 2) no bkpt +/- 100bp; N=~5.9k
 Black line: window smoothed (window_size=10, step=1)



Page 49: [26] Deleted Author 4/1/18 9:22:00 PM

[JZ2MG: special attention. To disc next week]

make a response for Orli using 4.9, the other referee thinks matching doesn't make sense

Page 49: [27] Deleted Author 4/1/18 9:22:00 PM


Apart from the advantage of single-cell analyses of enabling examination of complex cancer cell biology, there is, moreover, reason to believe that single-cell analyses may capture important tumor biology present *in vivo*. Cancers that result from a single progenitor cell, or homogenous progenitor population, provide a justification for the use of single-cell analyses and comparisons. There is evidence that a number of cancers may develop according to the cancer stem-cell model, which posits that it is only a small population of stem-like cells that are responsible for tumor development and observed intratumoral heterogeneity (PMID: 24607403). Understanding the biology of a single cells in the progenitor population may be sufficient to gain perspective on the tumor landscape as a whole.

Nonetheless, some of our analyses are should be particularly robust to the presence and activities of stromal and infiltrating cells. For example, our BMR calculations should not largely be affected by stromal tissue epigenetics, because clonally-amplified mutations detected by bulk sequencing will tend to accrue to a much greater extent in cells descendant from the cell-of-origin of the cancer cell much more so than associated normal tissue.

In addition, even when there is genomic heterogeneity observed across tumor clones and subclones, the main driver mutations and phenotypic traits may be widely shared among cells

(PMID: 3944607, 21376230). For example, in a single-cell sequencing analysis of colon cancer, the primary drivers TP53 and APC were present in the majority of cells across clones, with other mutations showing greater heterogeneity. (PMID: 24699064) Furthermore, even when there is substantial initial genomic and phenotypic heterogeneity, tumors may tend to converge to a genomic and phenotypic equilibrium (e.g. to a stem-like state) as has been shown in a number of studies on breast cancer tumor evolution (PMID: 21854987, 21498687, 22472879). As we have shown in the revised manuscript that, the conclusions we made from the cell lines correlate well with the observations from primary cancer patients.

We predicted the regulatory activities of transcription factor (TF) MYC using a ChIP-Seq profile in MCF-7 cells. We found that the MYC regulatory activity is highly correlated with the MYC expression across TCGA breast tumors (Supplementary Figure Xa). For most TFs, their regulatory activities predicted using ENCODE ChIP-Seq profile in cell lines are significantly correlated with their expression levels across breast tumors (Supplementary Figure Xb). Moreover, using the same MCF-7 ChIP-Seq profile, the MYC regulatory activity predicted for lung tumors is also significantly correlated with MYC expression level in TCGA lung cancer (Supplementary Figure Xa). These results indicate that the ChIP-Seq profiles from a particular cell line can capture regulatory targets in human tumors from diverse cancer types. To select ChIP-Seq or eCLIP profiles that are representative of the regulatory targets in human cancers, we only reported the results of TFs or RBPs whose regulatory activities are significantly correlated with their gene expression level in each TCGA cohort (Supplementary Figure Xc).



Supplementary Figure X. The clinical relevance of ENCODE cell line data in human primary tumors.

(a) The correlation between MYC expression level and regulatory activity across tumors. The MYC regulatory activity in each tumor was predicted using the ChIP-Seq profile in MCF-7 cell line. The Pearson correlation between MYC gene expression level and regulatory activity were computed across

tumors in each cancer type. The statistical significance of Pearson correlation was tested by the two-sided student t-test. BRCA: breast invasive carcinoma. LUSC: lung squamous carcinoma.

(b) The distribution of correlation p -values in TCGA breast cancer. For each TF, we tested the statistical significance of Pearson correlation between TF expression levels and regulatory activities predicted across tumors through two-sides student t tests as panel a. For TCGA breast cancer cohort, most p -values are very significant with a few non-significant values.

The fraction of regulators with statistically significant correlations in different cancer types for ChIP-Seq and eCLIP networks. In each TCGA cancer type, we computed the correlations between regulator expression levels and regulatory activities across tumors for all regulators (TFs, or RBPs). We selected regulators with statistically significant correlations through two-sided student t test (FDR < 0.05).

Page 58: [29] Deleted	Author	4/1/18 9:22:00 PM
-----------------------	--------	-------------------

Part of the previous

#####7mar:fight-outofscope

#####7mar - comparisons w/ other methods

###21mar - Inigo's paper is not about BMR/driver discovery

in response doc, praise referee, do analysis to compare Inigo's method

Page 58: [30] Deleted	Author	4/1/18 9:22:00 PM
-----------------------	--------	-------------------

Referee

Comment

Page 59: [31] Deleted	Author	4/1/18 9:22:00 PM
-----------------------	--------	-------------------

Author

Response


Page 59: [32] Moved to page 58 (Move #8)	Author	4/1/18 9:22:00 PM
--	--------	-------------------

In our initial submission, the assumption is that we were trimming off the nonfunctional sites while preserving the functional ones. Two examples can explain the motivation of this assumption.

1) Enhancers: Traditionally, enhancers were called as a 1kb peak regions, which admittedly introduced a lot of obviously nonfunctional sites. We believe we can get functional region more accurately by trimming the enhancers down using the exact shapes of many histone marks and further integration with STARR-seq and Hi-C data.

2) TFBS hotspots around the promoter region of WDR74. Instead of testing the conventional up to 2.5K promoter region, we can trim the test set to a core set of the promoter region where many TFs bind, which perfectly correlates with the mutation hotspots (red block) for this well-known driver site (blue line for pan-cancer and green line for liver cancer).

Page 59: [33] Deleted	Author	4/1/18 9:22:00 PM
-----------------------	--------	-------------------



Page 60: [34] Deleted **Author** **4/1/18 9:22:00 PM**

Following the reviewer’s suggestions, we show in a formal power analysis now in the supplement that the most important contribution to power comes from including additional functional sites, this is of course by the extended gene concept and then secondarily, from removing non-functional sites, but to a lesser extent. The assumption in our compacting annotations is that we can accurately distinguish the more important functional nucleotides from the less important ones through the guidance of many functional characterization assays.

However, we are admittedly making assumptions and the referee is completely correct in pointing this out. We have tried to be more precise in the text that we’re basically assuming that the large number of encode assays when integrated allows us to more directly get at the functionally important nucleotides, but this of course is an assumption. It’s hard to really tell to what degree one can success in finding the current events in cancer. It’s hard to back this up with the gold standard, but I think that some of the points are self evidently obvious. We’ve tried to make this clear in text and thank the referee for pointing this out.

Page 61: [35] Deleted **Author** **4/1/18 9:22:00 PM**

As for the enhancer part, with the ensemble method, for example, we can get more accurate annotation and pin-point to sequences where transcription factors would actually bind to. To estimate the false positive rate would not be very practical at this stage as there is no gold-standard experiment that could assert an predicted enhancer is definitely negative. Here we took the FANTOM enhancer data set and assess the overlap percentage of our enhancer annotation in each ensemble step. We show that each ensemble step indeed increases the percentage of overlap between our annotation and the FANTOM enhancer set. The overlap percentage for our annotation is much higher than that of the Roadmap annotation, and is also higher than the main encyclopedia enhancer annotation annotation (ccRE).

Page 65: [36] Moved to page 65 (Move #13) **Author** **4/1/18 9:22:00 PM**

<TYPE>\$\$\$BMR,\$\$\$Text
 <ASSIGN>@@@JZ
 <PLAN>&&&AgreeFix
 <STATUS>%%%

Page 65: [37] Deleted Author 4/1/18 9:22:00 PM

DONE

Page 65: [38] Moved to page 65 (Move #12) Author 4/1/18 9:22:00 PM

<TYPE>\$\$\$BMR,\$\$\$Text
<ASSIGN>@@@JZ
<PLAN>&&&AgreeFix
<STATUS>%%%100DONE

Page 66: [39] Moved to page 60 (Move #10) Author 4/1/18 9:22:00 PM

<ID>REF5.


Page 66: [40] Deleted Author 4/1/18 9:22:00 PM

14 – Power analysis for compact annotation

<TYPE>\$\$\$Power,\$\$\$Calc
<ASSIGN>@@@JZ
<PLAN>&&&AgreeFix
<STATUS>%%%100DONE

[JZ2MG: feel the three power related questions can be combined]


Referee Comment	How do the new "compact annotations" lead to improved results over traditional annotations?
Author Response	<p>We thank the referee for pointing this out. We have made it more clear in our supplementary file. In our initial submission, the assumption is that we were trimming off the nonfunctional sites while preserving the functional ones. Two examples can explain the motivation of this assumption.</p> <p>1) <u>Enhancers</u>: Traditionally, enhancers were called as a 1kb peak regions, which admittedly introduced a lot of obviously nonfunctional sites. We believe we can get functional region more accurately by trimming the enhancers down using the exact shapes of many histone marks and further integration with STARR-seq and Hi-C data.</p> <p>2) <u>TFBS hotspots around the promoter region of WDR74</u>. Instead of testing the conventional up to 2.5K promoter region, we can trim the test set to a core set of the promoter region where many TFs bind, which perfectly correlates with the mutation hotspots (red block) for this well-known driver site (blue line for pan-cancer and green line for liver cancer).</p>



Excerpt 3 From Revised Manuscript


Regarding quality of enhancers

As for the enhancer part, with the ensemble method, for example, we can get more accurate annotation and pin-point to sequences where transcription factors would actually bind to. To estimate the false positive rate would not be very practical at this stage as there is no gold-standard experiment that could assert an predicted enhancer is definitely negative. Here we took the FANTOM enhancer data set and assess the overlap percentage of our enhancer annotation in each ensemble step. We show that each ensemble step indeed increases the percentage of overlap between our annotation and the FANTOM enhancer set. The overlap percentage for our annotation is much higher than that of the Roadmap annotation, and is also higher than the main encyclopedia enhancer annotation annotation (ccRE).




Excerpt 4 From Revised Manuscript


Our original result




Result using alternative gene expression data from GEO







Our original result



Result using alternative gene expression data from GEO







We would like to clarify that the rewiring index is based on the fraction of regulatory edge changes between two cellular contexts. The rewiring index is also normalized across all regulatory proteins, and the sign

reflects the direction of rewiring. Details of rScore derivation can be found in Supplementary 5.3. Given this, we assume a null hypothesis to be no change in regulatory edge across cell types. We expect no or minimal change in edges when two cellular contexts are similar. To demonstrate, we selected all available GM12878 ChIP-seq experiments that have at least two replicates, and we performed the same rewiring analysis between isogenic replicates of the same cellular context. The edge changes between two networks will be simply a noise from ChIP-seq experiments.

As expected, when two cellular context are similar, as shown in “baseline”, minimal number of edges do change targets. However, in “rewiring”, TF do change targets extensively when compared across cancerous (K562) to normal (GM12878) cell lines.




p-value = 8.72e-17




Author Response

The eight regions were selected from our integrative promoter and enhancer



<Figure update candidate: Gene expression profiles of all available ENCODE RNA-seq experiments show that all stem-like cell types form a cluster (Blue). Gene expression quantifications were projected onto 2D space using reference component analysis.>



<Shadow figure of RCA>